



Physics

1. Find the acceleration of the two blocks. The system is initially at rest and the friction coefficient between block A and block B is $\mu = 0.5$. The ground below block B is smooth. A force $F = 80 \text{ N}$ is applied horizontally to block A. Both blocks have a mass of 10 kg . Take $g = 10 \text{ m/s}^2$.




- A) $a_A = 8 \text{ m/s}^2, a_B = 0 \text{ m/s}^2$ B) $a_A = a_B = 8 \text{ m/s}^2$
 C) $a_A = 6 \text{ m/s}^2, a_B = 2 \text{ m/s}^2$ D) $a_A = a_B = 4 \text{ m/s}^2$

2. Find the acceleration of the two blocks. The system is initially at rest and the friction coefficient between block A and block B is $\mu = 0.4$. The ground below block B is smooth. A force $F = 100 \text{ N}$ is applied horizontally to block A. Both blocks have a mass of 10 kg . Take $g = 10 \text{ m/s}^2$.



- A) $a_A = a_B = 5 \text{ m/s}^2$ B) $a_A = 10 \text{ m/s}^2, a_B = 0 \text{ m/s}^2$
 C) $a_A = 6 \text{ m/s}^2, a_B = 4 \text{ m/s}^2$ D) $a_A = a_B = 4 \text{ m/s}^2$


3. Find the acceleration of the two blocks. The system is initially at rest and the friction coefficient between block A and block B is $\mu = 0.5$. The ground below block B is smooth. A force $F = 120 \text{ N}$ is applied horizontally to block A. The mass of block A is 20 kg and the mass of



block B is 10 kg. Take $g = 10 \text{ m/s}^2$.


- A) $a_A = 12 \text{ m/s}^2, a_B = 0 \text{ m/s}^2$ B) $a_A = a_B = 6 \text{ m/s}^2$
 C) $a_A = a_B = 4 \text{ m/s}^2$ D) $a_A = 8 \text{ m/s}^2, a_B = 2 \text{ m/s}^2$

4. Initially the system is at rest. Find the minimum value of F for which sliding starts between the two blocks. The friction coefficient between block A and block B is $\mu = 0.5$ and the ground below block B is smooth. The mass of block A is 10 kg and the mass of block B is 30 kg. Take $g = 10 \text{ m/s}^2$.




- A) 150 N B) 200 N
 C) 250 N D) 100 N

5. In the given figure, find the accelerations and the friction forces involved for block A (4 kg) and block B (6 kg) when a force $F = 20 \text{ N}$ is applied to block A. The friction coefficient between A and B is $\mu = 0$ and between B and ground G is $\mu' = 0.5$. The system is initially at rest. Take $g = 10 \text{ m/s}^2$



- A) $a_A = 2 \text{ m/s}^2, a_B = 2 \text{ m/s}^2, f_{AB} = 12 \text{ N}, f_{BG} = 0 \text{ N}$ B) $a_A = 0 \text{ m/s}^2, a_B = 0 \text{ m/s}^2, f_{AB} = 20 \text{ N}, f_{BG} = 20 \text{ N}$
 C) $a_A = 5 \text{ m/s}^2, a_B = 0 \text{ m/s}^2, f_{AB} = 0 \text{ N}, f_{BG} = 5 \text{ N}$ D) $a_A = 0 \text{ m/s}^2, a_B = 2 \text{ m/s}^2, f_{AB} = 0 \text{ N}, f_{BG} = 5 \text{ N}$


6. In the given figure, find the accelerations and the friction forces involved for block A (6 kg) and block B (12 kg) when a force $F = 42 \text{ N}$ is applied to block A. The friction coefficient between A and B is $\mu = 0.5$ and between B and ground G is $\mu' = 0.5$. The system is initially at rest. Take



$g = 10 \text{ m/s}^2$.

- A) $a_A = 2 \text{ m/s}^2, a_B = 0 \text{ m/s}^2, f_{AB} = 30 \text{ N}, f_{BG} = 3 \text{ N}$ B) $a_A = 3 \text{ m/s}^2, a_B = 1 \text{ m/s}^2, f_{AB} = 24 \text{ N}, f_{BG} = 3 \text{ N}$
 C) $a_A = 2 \text{ m/s}^2, a_B = 2 \text{ m/s}^2, f_{AB} = 30 \text{ N}, f_{BG} = 7 \text{ N}$ D) $a_A = 0 \text{ m/s}^2, a_B = 0 \text{ m/s}^2, f_{AB} = 0 \text{ N}, f_{BG} = 3 \text{ N}$


7. In the given figure, find the accelerations and the friction forces involved for block A (5 kg) and block B (10 kg) when a horizontal force $F = 170 \text{ N}$ is applied to block B. The friction coefficient between A and B is $\mu = 0.6$ and between B and ground G is $\mu' = 0.4$. The system is initially at rest.



Take $g = 10 \text{ m/s}^2$.

- A) $a_A = 6 \text{ m/s}^2, a_B = 14 \text{ m/s}^2, f_{AB} = 30 \text{ N}, f_{BG} = 60 \text{ N}$
 B) $a_A = 6 \text{ m/s}^2, a_B = 8 \text{ m/s}^2, f_{AB} = 30 \text{ N}, f_{BG} = 60 \text{ N}$
 C) $a_A = 8 \text{ m/s}^2, a_B = 8 \text{ m/s}^2, f_{AB} = 40 \text{ N}, f_{BG} = 60 \text{ N}$
 D) $a_A = 0 \text{ m/s}^2, a_B = 11 \text{ m/s}^2, f_{AB} = 0 \text{ N}, f_{BG} = 60 \text{ N}$


8. In the given figure, find the accelerations and the friction forces involved for block A (4 kg) and block B (8 kg) when a force $F = 84 \text{ N}$ is applied to block B. The friction coefficient between A and B is $\mu = 0.5$ and between B and ground G is $\mu' = 0.5$. The system is initially at rest. Take



$g = 10 \text{ m/s}^2$.


- A) $a_A = 5 \text{ m/s}^2, a_B = 10 \text{ m/s}^2, f_{AB} = 20 \text{ N}, f_{BG} = 20 \text{ N}$
 B) $a_A = 2 \text{ m/s}^2, a_B = 0 \text{ m/s}^2, f_{AB} = 20 \text{ N}, f_{BG} = 20 \text{ N}$
 C) $a_A = 2 \text{ m/s}^2, a_B = 2 \text{ m/s}^2, f_{AB} = 8 \text{ N}, f_{BG} = 60 \text{ N}$
 D) $a_A = 0 \text{ m/s}^2, a_B = 0 \text{ m/s}^2, f_{AB} = 0 \text{ N}, f_{BG} = 0 \text{ N}$

9. Two blocks of masses 2 kg (upper) and 4 kg (lower) are placed on a horizontal rough surface. The coefficient of friction between the blocks is 0.5 and between the lower block and the ground is 0.2. A horizontal force of 30 N is applied on the lower block. Find the acceleration of the system. ($g = 10 \text{ m/s}^2$)




- A) 2 m/s^2
 B) 3 m/s^2
 C) 4 m/s^2
 D) 5 m/s^2

10. Two blocks of masses 3 kg (upper) and 7 kg (lower) are placed on a smooth horizontal surface. The coefficient of friction between the two blocks is 0.5. A horizontal force of 15 N is applied on the upper block. Find the acceleration of the lower block. ($g = 10 \text{ m/s}^2$)




- A) 1.5 m/s^2
 B) 2 m/s^2
 C) 3 m/s^2
 D) 4 m/s^2

11. Two blocks of masses 4 kg (upper) and 6 kg (lower) are placed on a rough horizontal surface. The coefficient of friction between the blocks is 0.4 and between the lower block and the ground is 0.3. A horizontal force of 40 N is applied on the upper block. Find the acceleration of the lower block. ($g = 10 \text{ m/s}^2$)




- A) 2 m/s^2 B) 3 m/s^2
C) 4 m/s^2 D) 0 m/s^2

12. Two blocks of masses 5 kg (upper) and 10 kg (lower) are placed on a rough horizontal surface. The coefficient of friction between the blocks is 0.2, and between the lower block and the ground is 0.1. A horizontal force of 50 N is applied on the lower block. Find the acceleration of the upper block. ($g = 10 \text{ m/s}^2$)




- A) 4 m/s^2 B) 3 m/s^2
C) 2 m/s^2 D) 5 m/s^2

13. Two blocks of masses 6 kg (upper) and 9 kg (lower) are placed on a smooth horizontal surface. The coefficient of friction between the blocks is 0.5. A horizontal force of 45 N is applied on the upper block. Find the acceleration of the system. ($g = 10 \text{ m/s}^2$)




- A) 2 m/s^2 B) 3 m/s^2
C) 4 m/s^2 D) 5 m/s^2

14. Two blocks of masses 2 kg and 3 kg are placed one over the other on a rough horizontal surface. The coefficient of friction between the blocks is 0.2 and the coefficient of friction between the lower block and the ground is 0.3. A horizontal force of 30 N is applied on the lower block as shown. Assume $g = 10 \text{ m/s}^2$. Find the acceleration of the blocks and the frictional force between the two blocks.




- A) $a_{2\text{kg}} = 2 \text{ m/s}^2, a_{3\text{kg}} = \frac{11}{3} \text{ m/s}^2, f = 4 \text{ N}$ B) $a_{2\text{kg}} = 3 \text{ m/s}^2, a_{3\text{kg}} = 3 \text{ m/s}^2, f = 6 \text{ N}$
C) $a_{2\text{kg}} = 2 \text{ m/s}^2, a_{3\text{kg}} = 5 \text{ m/s}^2, f = 4 \text{ N}$ D) $a_{2\text{kg}} = 0 \text{ m/s}^2, a_{3\text{kg}} = 5 \text{ m/s}^2, f = 0 \text{ N}$

15. Two blocks of masses 4 kg and 6 kg are placed one over the other on a smooth (horizontal) surface. The coefficient of friction between the blocks is 0.4. A horizontal force of 20 N is applied on the upper block as shown. Assume $g = 10 \text{ m/s}^2$. Find the acceleration of the blocks and the frictional force between the two blocks.




- A) $a_{4\text{kg}} = 2 \text{ m/s}^2, a_{6\text{kg}} = 2 \text{ m/s}^2, f = 12 \text{ N}$
 B) $a_{4\text{kg}} = 5 \text{ m/s}^2, a_{6\text{kg}} = 2.67 \text{ m/s}^2, f = 16 \text{ N}$
 C) $a_{4\text{kg}} = 2 \text{ m/s}^2, a_{6\text{kg}} = 2 \text{ m/s}^2, f = 16 \text{ N}$
 D) $a_{4\text{kg}} = 1 \text{ m/s}^2, a_{6\text{kg}} = 1 \text{ m/s}^2, f = 10 \text{ N}$

16. Two blocks of masses 3 kg and 7 kg are placed one over the other on a rough horizontal surface. The coefficient of friction between the blocks is 0.5 and the coefficient of friction between the lower block and the ground is 0.2. A horizontal force of 50 N is applied on the upper block as shown. Assume $g = 10 \text{ m/s}^2$. Find the acceleration of the blocks and the frictional force between the two blocks.




- A) $a_{3\text{kg}} = 5 \text{ m/s}^2, a_{7\text{kg}} = 3 \text{ m/s}^2, f = 15 \text{ N}$
 B) $a_{3\text{kg}} = \frac{35}{3} \text{ m/s}^2, a_{7\text{kg}} = 0 \text{ m/s}^2, f = 15 \text{ N}$
 C) $a_{3\text{kg}} = 10 \text{ m/s}^2, a_{7\text{kg}} = 2 \text{ m/s}^2, f = 20 \text{ N}$
 D) $a_{3\text{kg}} = \frac{35}{3} \text{ m/s}^2, a_{7\text{kg}} = 5 \text{ m/s}^2, f = 15 \text{ N}$

17. Two blocks of masses 5 kg and 10 kg are placed one over the other on a horizontal surface. The coefficient of friction between the blocks is 0.3 and the coefficient of friction between the lower block and the ground is 0.4. A horizontal force of 60 N is applied on the lower block as shown. Assume $g = 10 \text{ m/s}^2$. Find the acceleration of the blocks and the frictional force between the two blocks.




- A) $a_{5\text{kg}} = 0 \text{ m/s}^2, a_{10\text{kg}} = 0 \text{ m/s}^2, f = 0 \text{ N}$
 B) $a_{5\text{kg}} = 3 \text{ m/s}^2, a_{10\text{kg}} = 2 \text{ m/s}^2, f = 15 \text{ N}$
 C) $a_{5\text{kg}} = 2 \text{ m/s}^2, a_{10\text{kg}} = 2 \text{ m/s}^2, f = 10 \text{ N}$
 D) $a_{5\text{kg}} = 0 \text{ m/s}^2, a_{10\text{kg}} = 1 \text{ m/s}^2, f = 5 \text{ N}$

18. Two blocks of masses 6 kg and 4 kg are placed one over the other on a smooth horizontal surface. The coefficient of friction between the blocks is 0.5. A horizontal force of 40 N is applied on the upper block as shown. Assume $g = 10 \text{ m/s}^2$. Find the acceleration of the blocks and the frictional force between the two blocks.



- A) $a_{6\text{kg}} = 1.67 \text{ m/s}^2, a_{4\text{kg}} = 7.5 \text{ m/s}^2, f = 30 \text{ N}$
 B) $a_{6\text{kg}} = 6.67 \text{ m/s}^2, a_{4\text{kg}} = 0 \text{ m/s}^2, f = 0 \text{ N}$
 C) $a_{6\text{kg}} = 4 \text{ m/s}^2, a_{4\text{kg}} = 4 \text{ m/s}^2, f = 30 \text{ N}$
 D) $a_{6\text{kg}} = 4 \text{ m/s}^2, a_{4\text{kg}} = 4 \text{ m/s}^2, f = 16 \text{ N}$


19. Two blocks of masses 2 kg and 3 kg are placed one over the other on a rough horizontal surface. The coefficient of friction between the two blocks is 0.4 and the coefficient of friction between the lower block and the ground is 0.2. A horizontal force of 30 N is applied on the lower block as shown in the figure. Find the acceleration of the two blocks.



- A) 2 m/s^2
 C) 4 m/s^2


- B) 3 m/s^2
 D) 5 m/s^2

20. A block of mass 3 kg is placed on another block of mass 5 kg, which rests on a smooth (horizontal) surface. The coefficient of friction between the two blocks is 0.5. A horizontal force of 18 N is applied on the upper block. Find the acceleration of each block and the frictional force between the blocks. (Assume $g = 10 \text{ m/s}^2$)




- A) $a_{3\text{kg}} = 2.25 \text{ m/s}^2, a_{5\text{kg}} = 2.25 \text{ m/s}^2, f = 11.25 \text{ N}$
 B) $a_{3\text{kg}} = 11.25 \text{ m/s}^2, a_{5\text{kg}} = 3 \text{ m/s}^2, f = 15 \text{ N}$
 C) $a_{3\text{kg}} = 6 \text{ m/s}^2, a_{5\text{kg}} = 0 \text{ m/s}^2, f = 0 \text{ N}$
 D) $a_{3\text{kg}} = 3 \text{ m/s}^2, a_{5\text{kg}} = 3 \text{ m/s}^2, f = 15 \text{ N}$

21. A block of mass 4 kg is placed on another block of mass 6 kg. The coefficient of friction between the two blocks is 0.5 and the coefficient of friction between the lower block and the ground is 0.4. A horizontal force of 50 N is applied on the upper block as shown. Find the acceleration of the blocks and the frictional force acting between the blocks. (Assume $g = 10 \text{ m/s}^2$)




- A) $a_{4\text{kg}} = 12.5 \text{ m/s}^2, a_{6\text{kg}} = 0 \text{ m/s}^2, f = 0 \text{ N}$
 B) $a_{4\text{kg}} = 5 \text{ m/s}^2, a_{6\text{kg}} = 1 \text{ m/s}^2, f = 20 \text{ N}$
 C) $a_{4\text{kg}} = 7.5 \text{ m/s}^2, a_{6\text{kg}} = 2 \text{ m/s}^2, f = 40 \text{ N}$
 D) $a_{4\text{kg}} = 7.5 \text{ m/s}^2, a_{6\text{kg}} = 0 \text{ m/s}^2, f = 20 \text{ N}$

22. Two blocks of masses 1 kg and 4 kg are placed one over the other. The horizontal surface and the interface between the blocks are smooth (frictionless). A horizontal force of 20 N is applied on the lower block. Find the acceleration of each block and the normal force between the blocks. (Assume $g = 10 \text{ m/s}^2$)




- A) $a_{1\text{kg}} = 0 \text{ m/s}^2, a_{4\text{kg}} = 5 \text{ m/s}^2, N = 10 \text{ N}$
 B) $a_{1\text{kg}} = 4 \text{ m/s}^2, a_{4\text{kg}} = 4 \text{ m/s}^2, N = 10 \text{ N}$
 C) $a_{1\text{kg}} = 0 \text{ m/s}^2, a_{4\text{kg}} = 5 \text{ m/s}^2, N = 50 \text{ N}$
 D) $a_{1\text{kg}} = 5 \text{ m/s}^2, a_{4\text{kg}} = 5 \text{ m/s}^2, N = 10 \text{ N}$

23. Two blocks of masses 2 kg and 3 kg are placed one over the other on a rough horizontal surface. The coefficient of friction between the two blocks is 0.2 and the coefficient of friction between the lower block and the ground is 0.5. A horizontal force of 16 N is applied on the lower block. Find the acceleration of each block. (Assume $g = 10 \text{ m/s}^2$)




- A) $a_{2\text{kg}} = 1 \text{ m/s}^2, a_{3\text{kg}} = 1 \text{ m/s}^2$
 B) $a_{2\text{kg}} = 0 \text{ m/s}^2, a_{3\text{kg}} = 2 \text{ m/s}^2$
 C) $a_{2\text{kg}} = 0 \text{ m/s}^2, a_{3\text{kg}} = 0 \text{ m/s}^2$
 D) $a_{2\text{kg}} = 2 \text{ m/s}^2, a_{3\text{kg}} = 3 \text{ m/s}^2$

24. Two blocks of masses 2 kg and 3 kg are placed one over the other as shown in the figure. The coefficient of friction between the two blocks is 0.5 and the ground is smooth. A horizontal force of 20 N is applied on the upper block. Take $g = 10 \text{ m/s}^2$. Then the acceleration of each block is




- A) 4 m/s^2
 B) 5 m/s^2
 C) 6 m/s^2
 D) 10 m/s^2

25. Two blocks of masses 3 kg and 2 kg are placed one over the other as shown. The coefficient of friction between the two blocks is 0.4 and the ground is smooth. A horizontal force of 30 N is applied on the lower block. Take $g = 10 \text{ m/s}^2$. Then the acceleration of each block is




- A) 5 m/s^2
 B) 6 m/s^2
 C) 8 m/s^2
 D) 10 m/s^2

26. Two blocks of masses 1 kg and 4 kg are placed one over the other as shown. The coefficient of friction between the two blocks is 0.5 and the coefficient of friction between the lower block and the ground is 0.2. A horizontal force of 50 N is applied on the lower block. Take $g = 10 \text{ m/s}^2$. Then the acceleration of the system is




- A) 5 m/s^2
 B) 6 m/s^2
 C) 8 m/s^2
 D) 10 m/s^2

27. Two blocks of masses 4 kg and 6 kg are placed one over the other as shown. The coefficient of friction between the two blocks is 0.3 and the coefficient of friction between the lower block and the ground is 0.4. A horizontal force of 40 N is applied on the upper block. Take $g = 10 \text{ m/s}^2$. Then the acceleration of the upper block is




- A) 2 m/s^2
 B) 3 m/s^2
 C) 7 m/s^2
 D) 5 m/s^2

28. Two blocks of masses 2 kg and 3 kg are placed one over the other as shown. The coefficient of friction between the two blocks is 0.4 and the coefficient of friction between the lower block and the ground is 0.3. A horizontal force of 45 N is applied on the upper block. Take $g = 10 \text{ m/s}^2$. Then the acceleration of the system is




- A) 5 m/s^2
 B) 7.4 m/s^2
 C) 9 m/s^2
 D) 11 m/s^2

29. Two blocks of masses 2 kg and 3 kg are placed one over the other on a rough horizontal surface. The coefficient of static friction between the blocks is 0.5 and between the lower block and the ground is 0.2. A horizontal force F is applied on the upper block. Find the maximum value of F for which the two blocks move together without slipping. (Take $g = 10 \text{ m/s}^2$)




- A) 10 N
 B) 15 N
 C) 20 N
 D) 25 N

30. Two blocks of masses 4 kg and 6 kg are placed one over the other on a smooth horizontal surface (no friction with ground). The coefficient of static friction between the blocks is 0.4. A horizontal force of 40 N is applied on the lower block. Find the acceleration of the blocks and the frictional force between the blocks. (Assume $g = 10 \text{ m/s}^2$)




- A) $a = 4 \text{ m/s}^2, f = 16 \text{ N}$ B) $a = 4 \text{ m/s}^2, f = 24 \text{ N}$
 C) $a = 2.5 \text{ m/s}^2, f = 10 \text{ N}$ D) $a = 6.67 \text{ m/s}^2, f = 16 \text{ N}$

31. Two blocks of masses 5 kg and 10 kg are placed one over the other on a rough horizontal surface. The coefficient of static friction between the blocks is 0.3 and between the lower block and the ground is 0.4. A horizontal force F is applied on the lower block. What is the maximum value of F for which the upper block does not slip over the lower block? Also find the acceleration of the system at this force. (Assume $g = 10 \text{ m/s}^2$)




- A) $F = 105 \text{ N}, a = 3 \text{ m/s}^2$ B) $F = 90 \text{ N}, a = 2 \text{ m/s}^2$
 C) $F = 120 \text{ N}, a = 4 \text{ m/s}^2$ D) $F = 75 \text{ N}, a = 1 \text{ m/s}^2$

32. A block of mass 2 kg is placed on another block of mass 8 kg, which is kept on a rough horizontal surface. The coefficient of static friction between the blocks is 0.6 and between the lower block and the ground is 0.3. A horizontal force of 30 N is applied on the upper block. Determine (i) the acceleration of the blocks and (ii) the frictional force between the blocks. (Take $g = 10 \text{ m/s}^2$)



- A) $a_{2\text{kg}} = 9 \text{ m/s}^2, a_{8\text{kg}} = 0 \text{ m/s}^2, f = 12 \text{ N}$
 B) $a_{2\text{kg}} = 5 \text{ m/s}^2, a_{8\text{kg}} = 2 \text{ m/s}^2, f = 10 \text{ N}$
 C) $a_{2\text{kg}} = 15 \text{ m/s}^2, a_{8\text{kg}} = 0 \text{ m/s}^2, f = 0 \text{ N}$
 D) $a_{2\text{kg}} = 9 \text{ m/s}^2, a_{8\text{kg}} = 1 \text{ m/s}^2, f = 12 \text{ N}$

33. Two blocks of masses 3 kg and 7 kg are placed one over the other on a rough horizontal surface. The coefficient of static friction between the blocks is 0.2 and between the lower block and the ground is 0.5. A horizontal force of 16 N is applied on the lower block. Find (i) the acceleration of the blocks and (ii) the frictional force between the blocks. (Take $g = 10 \text{ m/s}^2$)




- A) $a = 0 \text{ m/s}^2, f = 0 \text{ N}$
 B) $a = 1 \text{ m/s}^2, f = 3 \text{ N}$
 C) $a = 2 \text{ m/s}^2, f = 6 \text{ N}$
 D) $a = 0 \text{ m/s}^2, f = 6 \text{ N}$

34. A block of 3 kg is stopped against a wall by applying a force F perpendicular to the wall. If $\mu = 0.6$ then minimum value of F will be (take $g = 10 \text{ m/s}^2$)
- A) 300 N
 B) 50 N
 C) 5 N
 D) 18 N

35. A block of 8 kg is initially at rest on a rough horizontal surface. The coefficient of static friction between the block and the surface is $\mu_s = 0.75$ and the coefficient of kinetic friction is $\mu_k = 0.50$. Taking $g = 10 \text{ m/s}^2$, the horizontal force required to just set the block in motion is
- A) 40 N
 B) 80 N
 C) 60 N
 D) 20 N


36. A block of mass 4 kg is kept on a rough floor. The coefficient of static friction between the block and the floor is 0.6. If a horizontal force F of 6 N is applied on the block as shown in the new diagram above, the frictional force between the block and the floor will be (take $g = 10 \text{ m/s}^2$)



- A) 24 N
 B) 12 N
 C) 40 N
 D) 6 N

37. If a ladder weighing 300 N is placed against a smooth vertical wall having coefficient of friction between it and floor is 0.4, then what is the maximum force of friction available at the point of contact between the ladder and the floor
- A) 75 N
 B) 300 N
 C) 120 N
 D) 30 N

45. A 10kg block is placed on a rough inclined plane of angle 30° and is connected by a light string passing over a smooth pulley to a hanging 10kg block. The coefficient of kinetic friction between the block and the incline is $\mu = 0.2$. Take $g = 10\text{m/s}^2$. Find the acceleration of the system and the frictional force on the block.



- A) $a = 1.63\text{m/s}^2$, $f = 17.3\text{N}$ B) $a = 2.0\text{m/s}^2$, $f = 20\text{N}$
 C) $a = 1.0\text{m/s}^2$, $f = 10\text{N}$ D) $a = 2.5\text{m/s}^2$, $f = 17.3\text{N}$

Chemistry

46. On complete combustion, 0.3 g of an organic compound gave 0.2 g of CO_2 and 0.1 g of H_2O . The percentage composition of carbon and hydrogen in the compound, respectively is:

- A) 15.02% and 3.70% B) 18.18% and 3.70%
 C) 15.02% and 4.07% D) 3.70% and 18.18%


47. 1 g of sodium hydroxide was treated with 25 mL of 0.75 M HCl solution, the mass of sodium hydroxide left unreacted is equal to:

- A) 750 mg B) 250 mg
 C) Zero mg D) 200 mg

48. A compound X contains 32% of A, 20% of B and remaining percentage of C. Then, the empirical formula of X is: (Given atomic masses of A = 64; B = 40; C = 32 u)


- A) A_2BC_2 B) ABC_3
 C) AB_2C_2 D) ABC_4

49. The right option for the mass of CO_2 produced by heating 20 g of 20% pure limestone is: (Atomic mass of Ca = 40)



- A) 1.76 g B) 2.64 g
 C) 1.32 g D) 1.12 g

50. What mass of 95% pure CaCO_3 will be required to neutralise 50 mL of 0.5 M HCl solution according to the following reaction?



- A) 1.25 g B) 1.32 g
 C) 3.65 g D) 9.50 g

51. An organic compound contains 78% (by wt.) carbon and remaining percentage of hydrogen. The right option for the empirical formula of this compound is: [Atomic wt. of C is 12, H is 1]

- A) CH B) CH_2
 C) CH_3 D) CH_4

52. The number of moles of hydrogen molecules required to produce 20 moles of ammonia through Haber's process is:

- A) 40 B) 10
 C) 20 D) 30

53. A mixture of 2.3 g formic acid and 4.5 g oxalic acid is treated with conc. H_2SO_4 . The evolved gaseous mixture is passed through KOH pellets. Weight (in g) of the remaining product at STP will be:

- A) 1.4 g B) 3.0 g
 C) 2.8 g D) 4.4 g

54. What is the mass of the precipitate formed when 50 mL of 16.9% solution of AgNO_3 is mixed with 50 mL of 5.8% NaCl solution? (Ag = 107.8, N = 14, O = 16, Na = 23, Cl = 35.5)

- A) 3.5 g B) 7 g
 C) 14 g D) 28 g

55. 1.0 g of magnesium is burnt with 0.56 g O_2 in a closed vessel. Which reactant is left in excess and how much? (At. wt. Mg = 24, O = 16)

- A) Mg, 0.16 g B) O_2 , 0.16 g
C) Mg, 0.44 g D) O_2 , 0.28 g

56. When 22.4 liters of $H_{2(g)}$ is mixed with 11.2 liters of $Cl_{2(g)}$, each at S.T.P, the moles of $HCl_{(g)}$ formed is equal to:

- A) 1 mol of $HCl_{(g)}$ B) 2 mol of $HCl_{(g)}$
C) 0.5 mol of $HCl_{(g)}$ D) 1.5 mol of $HCl_{(g)}$

57. In an experiment it showed that 10 mL of 0.05 M solution of chloride required 10 mL of 0.1 M solution of $AgNO_3$, which of the following will be the formula of the chloride (X stands for the symbol of the element other than chlorine)?

- A) X_2Cl_2 B) XCl_2
C) XCl_4 D) X_2Cl

58. 25.3 g of sodium carbonate Na_2CO_3 is dissolved in enough water to make 250 mL of solution. If sodium carbonate dissociates completely, molar concentration of sodium ion, Na^+ and carbonate ions, CO_3^{2-} are respectively: (Molar mass of Na_2CO_3 = 106 g mol^{-1})

- A) 0.955 M and 1.910 M B) 1.910 M and 0.955 M
C) 1.90 M and 1.910 M D) 0.955 M and 0.955 M

59. 10 g of hydrogen and 64 g of oxygen were filled in a steel vessel and exploded. Amount of water produced in this reaction will be:

- A) 3 mol B) 4 mol
C) 1 mol D) 2 mol

60. An organic compound contains carbon, hydrogen and oxygen. Its elemental analysis gave C, 38.71% and H, 9.67%. The empirical formula of the compound would be:

- A) CHO B) CH_4O
C) CH_3O D) CH_2O

61. How many moles of lead (II) chloride will be formed from a reaction between 6.5 g of PbO and 3.2 g HCl?

- A) 0.011 B) 0.029
C) 0.044 D) 0.333

62. What volume of oxygen gas (O_2) measured at $0^\circ C$ and 1 atm, is needed to burn completely 1 L propane gas (C_3H_8) measured under the same conditions?

- A) 5 L B) 10 L
C) 7 L D) 6 L

63. An element X has the following isotopic composition: ^{200}X : 90%, ^{199}X : 8.0%, ^{202}X : 2.0%. The weighted average atomic mass of the naturally occurring element X is closest to:

- A) 201 amu B) 202 amu
C) 199 amu D) 200 amu

64. Volume of CO_2 obtained by the complete decomposition of 9.85 g of $BaCO_3$ is:

- A) 2.24 L B) 1.12 L
C) 0.84 L D) 0.56 L

65. In the reaction, $4NH_{3(g)} + 5O_{2(g)} \rightarrow 4NO_{(g)} + 6H_2O_{(l)}$, when 1 mole of ammonia and 1 mole of O_2 are made to react to completion:

- A) all the oxygen will be consumed B) 1.0 mole of NO will be produced
C) 1.0 mole of H_2O is produced D) all the ammonia will be consumed

66. The amount of zinc required to produce 224 mL of H_2 at STP on treatment with dilute H_2SO_4 will be:

- A) 65 g B) 0.065 g
C) 0.65 g D) 6.5 g

67. Mass of methane required to produce 22 g of CO_2 after complete combustion is: (Given molar mass in g mol^{-1} : C = 12.0, H = 1.0, O = 16.0)

- A) 8 g B) 16 g
C) 22 g D) 44 g

68. Number of moles of methane required to produce 22 g $CO_2(g)$ after combustion is $x \times 10^{-2}$ moles. The value of x is:

- A) 50
C) 100
B) 25
D) 10

69. A sample of $CaCO_3$ and $MgCO_3$ weighed 2.21 g is ignited to constant weight of 1.152 g. The composition of the mixture is: (Given molar mass in $g\ mol^{-1}$: $CaCO_3$: 100, $MgCO_3$: 84)

- A) 1.187 g $CaCO_3$ + 1.023 g $MgCO_3$
C) 1.187 g $CaCO_3$ + 1.187 g $MgCO_3$
B) 1.023 g $CaCO_3$ + 1.023 g $MgCO_3$
D) 1.023 g $CaCO_3$ + 1.187 g $MgCO_3$

70. Consider the reaction:

$3PbCl_2 + 2(NH_4)_3PO_4 \rightarrow Pb_3(PO_4)_2 + 6NH_4Cl$. If 72 mmol of $PbCl_2$ is mixed with 50 mmol of $(NH_4)_3PO_4$, then the amount of $Pb_3(PO_4)_2$ formed is:

- A) 24 mmol
C) 18 mmol
B) 36 mmol
D) 50 mmol

71. 10 mL of gaseous hydrocarbon on combustion gives 40 mL of $CO_2(g)$ and 50 mL of water vapour. Total number of carbon and hydrogen atoms in the hydrocarbon is:

- A) 14
C) 12
B) 10
D) 8

72. When $Fe_{0.93}O$ is heated in presence of oxygen, it converts to Fe_2O_3 . How many statements are correct among the following? A. The equivalent weight of $Fe_{0.93}O$ is $\frac{\text{Molecular weight}}{0.79}$. B. The number of moles of Fe^{2+} and Fe^{3+} in 1 mole of $Fe_{0.93}O$ is 0.79 and 0.14 respectively. C. $Fe_{0.93}O$ is metal deficient with lattice comprising of cubic closed packed arrangement of O^{2-} ions. D. The % composition of Fe^{2+} and Fe^{3+} in $Fe_{0.93}O$ is 85% and 15% respectively.

- A) 4
C) 2
B) 3
D) 1

73. What is the mass ratio of ethylene glycol ($C_2H_6O_2$, molar mass = $62\ g\ mol^{-1}$) required for making 500 g of 0.25 molal aqueous solution and 250 mL of 0.25 molar aqueous solution?

- A) 1 : 1
C) 2 : 1
B) 3 : 1
D) 1 : 2

74. Number of hydrogen atoms per molecule of a hydrocarbon A having molar mass $84\ g\ mol^{-1}$ and containing 85.8% carbon by mass is:

- A) 12
C) 8
B) 6
D) 10

75. When a hydrocarbon A undergoes combustion in the presence of air, it requires 9.5 equivalents of oxygen and produces 3 equivalents of water. What is the molecular formula of A?

- A) C_8H_6
C) C_6H_6
B) C_9H_9
D) C_9H_8

76. On complete combustion, 0.492 g of an organic compound gave 0.792 g of CO_2 . The % of carbon in the organic compound is:

- A) 44
C) 88
B) 22
D) 50

77. Zinc reacts with hydrochloric acid to give hydrogen and zinc chloride. The volume of hydrogen gas produced at STP from the reaction of 11.5 g of zinc with excess HCl is: (Given: Molar mass of Zn = $65.4\ g\ mol^{-1}$, Molar volume of H_2 at STP = $22.7\ L$)

- A) 4 L
C) 11.2 L
B) 2.24 L
D) 8 L

78. Assume carbon burns according to following equation:

$2C_{(s)} + O_{2(g)} \rightarrow 2CO_{(g)}$ When 12g carbon is burnt in 48g of oxygen, the volume of carbon monoxide produced is $____ \times 10^{-1}\ L$ at STP [nearest integer] [Given: Assume CO as ideal gas, Mass of C is $12\ g\ mol^{-1}$, Mass of O is $16\ g\ mol^{-1}$ and molar volume of an ideal gas at STP is $22.7\ L\ mol^{-1}$]

- A) 227
C) 448
B) 112
D) 224

79. If 5 moles of $BaCl_2$ is mixed with 2 moles of Na_3PO_4 , the maximum number of moles of $Ba_3(PO_4)_2$ formed is:

- A) 1 B) 2
C) 3 D) 5

80. 0.5 g of an organic compound (X) with 60% carbon will produce $x \times 10^{-1}$ g of CO_2 on complete combustion. The value of x is:

- A) 11 B) 22
C) 44 D) 5.5

81. A solution of sugar is obtained by mixing 200 g of its 25% solution and 500 g of its 40% solution (both by mass). The mass percentage of the resulting sugar solution is:

- A) 36% B) 25%
C) 40% D) 30%

82. An organic compound gives 0.220 g of CO_2 and 0.126 g of H_2O on complete combustion. If the % of carbon is 24 then the % of hydrogen is $x \times 10^{-1}$. The value of x is:

- A) 56 B) 28
C) 14 D) 70

83. 1 g of a carbonate (M_2CO_3) on treatment with excess HCl produces 0.01 mol of CO_2 . The molar mass of M_2CO_3 is:

- A) 100 g mol⁻¹ B) 84 g mol⁻¹
C) 106 g mol⁻¹ D) 120 g mol⁻¹

84. Compound A contains 8.7% Hydrogen, 74% Carbon and 17.3% Nitrogen. The molecular formula of the compound is, Given: Atomic masses of C, H and N are 12, 1 and 14 amu respectively. The molar mass of the compound A is 162, g mol⁻¹.

- A) $C_4H_6N_2$ B) C_2H_3N
C) C_5H_7N D) $C_{10}H_{14}N_2$

85. The complete combustion of 0.492 g of an organic compound containing C, H and O gives 0.7938 g of CO_2 and 0.4428 g of H_2O . The percentage of oxygen composition is:

- A) 46% B) 43.5%
C) 9.9% D) 53.4%

86. Production of iron in blast furnace follows

$Fe_3O_4(s) + 4CO(g) \rightarrow 3Fe(l) + 4CO_2(g)$. When 4.640 kg of Fe_3O_4 and 2.520 kg of CO react, the amount of iron produced is:

- A) 2200 g B) 1400 g
C) 3360 g D) 4640 g

87. 120 g of an organic compound containing C and H gives 330 g of CO_2 and 270 g of water on complete combustion. The percentage of carbon and hydrogen are:

- A) 40 and 60 B) 25 and 75
C) 60 and 40 D) 75 and 25

88. On complete combustion 0.30 g of an organic compound gave 0.20 g of carbon dioxide and 0.10 g of water. The percentage of carbon in the given organic compound is ____:

- A) 18% B) 24%
C) 12% D) 36%

89. The moles of methane required to produce 81 g of water after complete combustion is $x \times 10^{-2}$ mol. The value of x is:

- A) 225 B) 450
C) 112 D) 250

90. 116g of a substance upon dissociation reaction, yields 7.5g of hydrogen, 60g of oxygen and 48.5g of carbon. Given that the atomic masses of H, O and C are 1, 16 and 12 respectively. The data agrees with how many formulae of the following? A. CH_3COOH B. $HCHO$ C. CH_3OCH_3 D. CH_3CHO

- A) 2 B) 1
C) 3 D) 4

Botany

91. Vexillary aestivation and a diadelphous androecium are characteristic features of which family?

- A) brinjal B) sem
C) marigold D) mustard

92. Epipetalous stamens and a persistent calyx (often found in brinjal and tomato) are traits of:

- A) Malvaceae B) Solanaceae
C) Liliaceae D) Poaceae

93. The typical fruit type associated with mustard and radish is a:

- A) Caryopsis B) Siliqua
C) Cypsela D) Berry

94. A capitulum (head) inflorescence with ray and disc florets and a modified hairy calyx called a pappus belongs to:

- A) Poaceae B) Liliaceae
C) Asteraceae D) Fabaceae

95. The fruit type in cereals like wheat and maize where the pericarp is completely fused with the seed coat is a:

- A) Cypsela B) Caryopsis
C) Cremocarp D) Nut


96. Trimerous flowers, epiphyllous stamens, and a perianth composed of 6 tepals arranged in two whorls (3+3) are diagnostic features of:

- A) Brassicaceae B) Liliaceae
C) Fabaceae D) Solanaceae

97. Colchicine, an alkaloid used to induce polyploidy, is extracted from a plant belonging to which family?

- A) Solanaceae B) Liliaceae
C) Asteraceae D) Poaceae

98. Which of the following is the correct floral formula for the floral diagram



given below?

- | | |
|---|---|
| <p>A) $\oplus \ \text{♀} \ K_{2+2} \ C_4 \ A_{2+4} \ \underline{G}_{(2)}$</p> <p>C) $\oplus \ \text{♀} \ K_{(5)} \ C_{(5)} \ A_5 \ \underline{G}_{(2)}$</p> | <p>B) $\% \ \text{♀} \ K_{(5)} \ C_{1+2+(2)} \ A_{(9)+1} \ \underline{G}_1$</p> <p>D) $\text{Br} \ \oplus \ \text{♀} \ P_{(3+3)} \ A_{3+3} \ G_{(3)}$</p> |
|---|---|

99. Pulse-yielding plants like Gram, Arhar, Sem, Moong, and Soyabean belong to Fabaceae. Which of the following dye-yielding plants also belongs to this family?

- A) Indigofera B) Colchicum
C) Atropa D) Nicotiana

100.

Non-endospermic (exalbuminous) seeds are typically found in the mature seeds of:

- A) Solanaceae B) Liliaceae
C) Fabaceae D) Poaceae

101.

Epipetalous stamens, where the filaments are fused directly to the petals, are characteristic of:

- A) Solanaceae B) Liliaceae
C) Fabaceae D) Brassicaceae

102.

Tobacco (*Nicotiana tabacum*) and Petunia belong to which family?

- | | |
|----------------------|----------------------|
| A) Malvaceae | B) Solanaceae |
| C) Asteraceae | D) Poaceae |

103.

The ovary is unilocular initially but becomes bilocular due to the formation of a false septum called the replum. This phenomenon occurs in:

- | | |
|------------------------------|------------------------------------|
| A) Mustard and Radish | B) Pea and Bean |
| C) Tomato and Onion | D) Cotton and Lady's finger |

104.

Monadelphous condition where all filaments fuse together to form a long staminal tube enclosing the style is found in:

- | | |
|-----------------|----------------------|
| A) Pea | B) China rose |
| C) Lemon | D) Sunflower |

105.

An outer whorl of bracteoles resembling an extra calyx layer, called the epicalyx, is missing in Solanaceae but universally present in:

- | | |
|---------------------|------------------------|
| A) Liliaceae | B) Poaceae |
| C) Malvaceae | D) Brassicaceae |

106.

Syngenesious stamens (anthers are fused into a tube around the style while the filaments remain entirely free) are seen in:

- | | |
|--------------------|----------------------------------|
| A) Solanum | B) Helianthus (Sunflower) |
| C) Brassica | D) Hibiscus |

107.

An inferior, unilocular ovary with a single basal ovule that ripens into a cypsela fruit is typical of:

- | | |
|----------------------|--------------------|
| A) Liliaceae | B) Poaceae |
| C) Asteraceae | D) Fabaceae |

108.

Which of the following is incorrect regarding the family Fabaceae?

- | | |
|---|----------------------------------|
| A) Monocarpellary superior ovary | B) Legume or pod fruit |
| C) Non-endospermic seeds | D) Actinomorphic symmetry |

109.

An obligately placed, bicarpellary, syncarpous superior ovary with an obliquely swollen placenta and axile placentation is found in:

- | | |
|--------------------|--------------------|
| A) Petunia | B) Aloe |
| C) Brassica | D) Sesbania |

110.

Tetradynamous stamens (2 short + 4 long) and a false septum called a replum are found in:

- | | |
|------------------------|--------------------|
| A) Malvaceae | B) Fabaceae |
| C) Brassicaceae | D) Poaceae |

111.

Monadelphous stamens forming a staminal tube around the style and a 3 to many-lobed epicalyx are unique landmarks of:

- | | |
|----------------------|---------------------|
| A) Solanaceae | B) Malvaceae |
| C) Asteraceae | D) Liliaceae |

112.

Which of the following sets of plants entirely belongs to Malvaceae?

- | | |
|---|-----------------------------------|
| A) Cotton, China rose, Lady's finger | B) Tomato, Brinjal, Potato |
| C) Sunflower, Marigold, Lettuce | D) Wheat, Maize, Rice |

113.

Syngenesious stamens (anthers fused, filaments free) and an inferior ovary are key traits of:

- | | |
|----------------------|------------------------|
| A) Asteraceae | B) Solanaceae |
| C) Malvaceae | D) Brassicaceae |

114.

Scale-like structures called lodicules, which represent a reduced perianth, are found in:

- | | |
|--------------|--------------|
| A) Liliaceae | B) Poaceae |
| C) Fabaceae | D) Malvaceae |

115.

Trimerous flowers, epiphyllous stamens, and a perianth composed of 6 tepals arranged in two whorls (3+3) are diagnostic features of:

- | | |
|-----------------|---------------|
| A) Brassicaceae | B) Liliaceae |
| C) Fabaceae | D) Solanaceae |

116.

Colchicine, an alkaloid used to induce polyploidy, is extracted from a plant belonging to which family?

- | | |
|---------------|--------------|
| A) Solanaceae | B) Liliaceae |
| C) Asteraceae | D) Poaceae |

117.

Diadelphous stamens [(9) + 1] and vexillary (descendingly imbricate) aestivation of the corolla are the signature features of:

- | | |
|---------------|-----------------|
| A) Solanaceae | B) Brassicaceae |
| C) Fabaceae | D) Liliaceae |

118.

The calyx in Asteraceae is modified into hairy, feathery, or scale-like structures that help in wind-dispersal of seeds. This modified structure is known as

- | | |
|-------------|-----------|
| A) Lodicule | B) Pappus |
| C) Epicalyx | D) Glume |

119.

An obliquely placed, bicarpellary, syncarpous superior ovary with an extremely swollen placenta and numerous ovules is found in:

- | | |
|-------------|------------|
| A) Brassica | B) Solanum |
| C) Pisum | D) Allium |

120.

Which of the following medicinal plants is a member of the family Solanaceae?

- | | |
|---------------------|-------------------------------------|
| A) Aloe vera | B) Withania somnifera (Ashwagandha) |
| C) Gloriosa superba | D) Indigofera tinctoria |

121.

Tobacco (*Nicotiana tabacum*) and Petunia belong to which family?

- | | |
|---------------|---------------|
| A) Malvaceae | B) Solanaceae |
| C) Asteraceae | D) Poaceae |

122.

Tetradynamous androecium (six stamens arranged in two whorls: 2 short outer and 4 long inner) is the characteristic feature of:

- | | |
|-----------------|---------------|
| A) Malvaceae | B) Solanaceae |
| C) Brassicaceae | D) Liliaceae |

123.

Cruciform corolla (four free petals arranged in the form of a cross) is found in:

- | | |
|------------------------|---------------------------|
| A) Brassica campestris | B) Hibiscus rosa-sinensis |
| C) Solanum nigrum | D) Helianthus annuus |

124.

Monothealous, kidney-shaped (reniform) anthers are structurally unique to which family?

- | | |
|---------------|---------------|
| A) Solanaceae | B) Malvaceae |
| C) Fabaceae | D) Asteraceae |

125.

Cotton (*Gossypium*), Lady's Finger (*Abelmoschus esculentus*), and Mallow belong to:

- | | |
|-----------------|--------------|
| A) Brassicaceae | B) Malvaceae |
| C) Fabaceae | D) Poaceae |

126.

An inferior, unilocular ovary with a single basal ovule that ripens into a cypsela fruit is typical of:

- A) Liliaceae**
- B) Poaceae**
- C) Asteraceae**
- D) Fabaceae**

127.

The highly reduced, scale-like perianth structures in the flowers of grasses that swell up to force the glumes apart are called:

- A) Lodicules**
- B) Pappus**
- C) Lemmas**
- D) Paleas**

128.

A characteristic monocotyledonous floral feature consisting of a perianth made of 6 tepals arranged in two whorls (3+3), often showing epiphyllous stamens, belongs to:

- A) Fabaceae**
- B) Liliaceae**
- C) Solanaceae**
- D) Asteraceae**

129.

Tricarpellary, syncarpous, superior ovary with axile placentation and trimerous flowers are diagnostic parameters for:

- A) Brassicaceae**
- B) Solanaceae**
- C) Liliaceae**
- D) Gramineae**

130.

Onion (*Allium cepa*) and Garlic (*Allium sativum*) modify their leaves into underground fleshy scales for food storage. They belong to:

- A) Solanaceae**
- B) Liliaceae**
- C) Poaceae**
- D) Malvaceae**

131.

Pulses like Gram (chana), Arhar (tur), Sem, Moong, and Masoor are elite protein sources belonging to:

- A) Solanaceae**
- B) Fabaceae**
- C) Liliaceae**
- D) Asteraceae**

132.

Which of the following is an example of an ornamental plant belonging to the monocot family Liliaceae?

- A) Tulipa (Tulip) and Gloriosa (Glory lily)** **B) Petunia**
C) Lupinus **D) Sweet pea**

133.

Asparagus, widely used as a gourmet vegetable and a traditional Ayurvedic medicine (Shatavari), belongs to:

- A) Fabaceae**
- B) Liliaceae**
- C) Solanaceae**
- D) Asteraceae**

134.

Match the following plant examples with their correct families:

Column I	Column II
A. Petunia	1. Poaceae
B. Triticum	2. Solanaceae
C. Aloe	3. Liliaceae
D. Pisum	4. Fabaceae

- A) A**
- B) B**
- C) C**
- D) D**

135.

Identify the correct statement regarding the aestivation of calyx in Brassicaceae:

- A) Valvate**
- B) Imbricate**
- C) Vexillary**
- D) Twisted**

Zoology

136.

In an ecosystem, the pyramid of energy is:

- A) Always upright.
- B) Sometimes inverted
- C) Always inverted
- D) Both upright and inverted

137.

The pyramid of biomass in a marine ecosystem is usually:

- A) Upright
- B) Inverted
- C) Linear
- D) Circular

138.

Which of the following best describes a food chain?

- A) It represents different populations living together.
- B) It shows the unidirectional flow of energy from producers to top consumers.
- C) It indicates only predator-prey relations.
- D) It shows population growth in an ecosystem.

139.

In forest ecosystems, the pyramid of numbers is generally:

- A) Circular
- B) Elliptical
- C) Upright
- D) Inverted

140.

The correct sequence in a typical food chain is:

- A) Herbivore → Producer → Carnivore
- B) Producer → Herbivore → Carnivore
- C) Carnivore → Herbivore → Producer
- D) Producer → Carnivore → Herbivore

141.

Match Column I (Ecosystem type) with Column II (Pyramid structure):

Column I	Column II
A) Marine	1. Upright pyramid of biomass
B) Forest	2. Inverted pyramid of biomass
C) Grassland	3. Pyramid of energy
D) Pond	4. Upright pyramid of number

- A) A-2, B-1, C-3, D-4
- B) A-2, B-1, C-1, D-2
- C) A-1, B-2, C-3, D-4
- D) A-3, B-4, C-1, D-2

142.

Match the following trophic levels to their examples:

Column I	Column II
A) Producer	1. Plants
B) Primary consumer	2. Cow
C) Secondary consumer	3. Frog
D) Tertiary consumer	4. Snake

- A) A-1, B-2, C-3, D-4
- B) A-2, B-1, C-3, D-4
- C) A-1, B-2, C-4, D-3
- D) A-1, B-4, C-2, D-3

143.

In the following, upright pyramid of number, the base represents:

- A) Top carnivores
- B) Primary producers
- C) Herbivores
- D) Secondary consumers

144.

Which statement is correct? With respect to Energy pyramid for a terrestrial food chain,

- A) Energy available at the top is more than at the base.
- B) Most energy is lost as heat at each transfer.
- C) Consumers are always at the base.
- D) Producers receive energy from both consumers and decomposers.

145.

Refer to the sequence: Grass → Grasshopper → Mouse → Owl. The mouse represents which trophic level?

- A) Primary consumer
- B) Secondary consumer
- C) Tertiary consumer
- D) Producer

146.

An ecosystem has a GPP of 1000 kcal/m²/year. Plants lose 600 kcal/m²/year via respiration. What is its NPP?

- A) 400 kcal/m²/year
- B) 1000 kcal/m²/year
- C) 1600 kcal/m²/year
- D) 600 kcal/m²/year

147.

Statement I: Decomposition is the process by which detritus is broken down into simpler substances by microbes. Statement II: The rate of decomposition is faster if detritus is rich in lignin and chitin.

- A) Both I and II correct
- B) Both I and II incorrect
- C) I correct, II incorrect
- D) I incorrect, II correct

148.

Detritivores contribute to decomposition by:

- A) Fragmenting detritus into smaller pieces
- B) Directly mineralizing organic matter
- C) Producing humus from detritus
- D) Photosynthesizing organic compounds

149.

Which of the following is NOT a decomposer?

- A) Fungi
- B) Earthworm
- C) Bacteria
- D) Green plants

150.

Which factor does NOT affect the rate of decomposition?

- A) Temperature
- B) Humidity
- C) Amount of sunlight
- D) Chemical composition of detritus

151.

Decomposition results in the conversion of:

- A) Organic matter into inorganic nutrients
- B) Inorganic nutrients into organic compounds
- C) Biomass into energy directly
- D) Detritus into living plants

152.

Humification is:

- A) Breakdown of detritus into smaller particles
- B) Formation of dark-colored amorphous substance from organic matter
- C) Photosynthesis by decomposers
- D) Uptake of nutrients by plants

153.

Which of these is a product of decomposition?

- A) Oxygen
- B) Carbon dioxide
- C) Glucose
- D) Nitrogen gas

154.

In ecosystems, decomposers mainly help in:

- A) Energy flow
- B) Nutrient cycling
- C) Primary productivity
- D) Biomass accumulation

155.

The process of detritus breakdown by detritivores is called:

- A) Catabolism
- B) Humification
- C) Fragmentation
- D) Nitrification

156.

Which of the following statements about lignin is true?

- A) It enhances the rate of decomposition
- B) It slows down the rate of decomposition
- C) It has no effect on decomposition
- D) It is produced by decomposers

157.

The major decomposers in an ecosystem are:

- A) Earthworms and fungi
B) Fungi and bacteria
C) Earthworms and protozoa
D) Bacteria and arthropods

158.

When temperature increases, the rate of decomposition usually:

- A) Decreases
B) Increases
C) Remains constant
D) First decreases then increases

159.

Statement I: Humus is resistant to microbial decomposition and provides nutrients to soil. Statement II: Humus is highly soluble in water.

- A) Both true
B) Both false
C) I true, II false
D) I false, II true

160.

During decomposition, bacteria mainly help in:

- A) Mineralization of organic matter
B) Fixing atmospheric nitrogen
C) Photosynthesis
D) Formation of detritus

161.

The rate of decomposition is faster in:

- A) Cold and dry conditions
B) Cold and wet conditions
C) Warm and moist conditions
D) Hot and dry conditions

162.

Match the following:

Column A (Concept)	Column B (Description)
(a) Gross Primary Productivity (GPP)	(1) Energy used by plants in respiration
(b) Net Primary Productivity (NPP)	(2) Total solar energy captured by plants
(c) Respiration	(3) Energy left after respiration for growth
(d) Energy flow	(4) Unidirectional transfer of energy in ecosystem

- A) 2, (b) 3, (c) 1, (d) 4
B) 3, (b) 2, (c) 4, (d) 1
C) 1, (b) 4, (c) 2, (d) 3
D) 4, (b) 1, (c) 3, (d) 2

163.

Match the following organisms with their trophic levels:

Column A (Organism)	Column B (Trophic Level)
(a) Grass	(1) Secondary consumer
(b) Frog	(2) Producer
(c) Snake	(3) Primary consumer
(d) Grasshopper	(4) Tertiary consumer

- A) 2, (b) 3, (c) 4, (d) 1
B) 3, (b) 2, (c) 1, (d) 4
C) 2, (b) 1, (c) 4, (d) 3
D) 4, (b) 3, (c) 2, (d) 1

164.

Match the following types of ecological pyramids with their characteristic:

Column A	Column B
(a) Pyramid of numbers	(1) Always upright
(b) Pyramid of biomass	(2) Can be inverted in aquatic food chain
(c) Pyramid of energy	(3) Represents number of organisms
(d) Pyramid of productivity	(4) Always upright and shows energy flow

- A) 3, (b) 2, (c) 4, (d) 1
B) 4, (b) 3, (c) 1, (d) 2
C) 4, (b) 1, (c) 3, (d) 2
D) 3, (b) 4, (c) 2, (d) 1

165.

Match the following types of ecosystems with example:

Column A (Ecosystem)	Column B (Example)
(a) Terrestrial	(1) Pond
(b) Aquatic	(2) Grassland
(c) Freshwater ecosystem	(3) Desert
(d) Desert	(4) Lake

- A) 2, (b) 1, (c) 4, (d) 3
B) 3, (b) 2, (c) 1, (d) 4
C) 1, (b) 3, (c) 2, (d) 4
D) 4, (b) 1, (c) 3, (d) 2

166.

Match the following terms with their definitions:

Column A (Term)	Column B (Definition)
(a) Detritus	(1) Dead organic matter
(b) Producers	(2) Organisms that make food using sunlight
(c) Decomposers	(3) Organisms breaking down organic wastes
(d) Consumers	(4) Organisms that feed on other organisms

A) 1, (b) 2, (c) 3, (d) 4

B) 2, (b) 3, (c) 4, (d) 1

C) 3, (b) 4, (c) 2, (d) 1

D) 4, (b) 1, (c) 3, (d) 2

167.

Match the following processes with associated terms:

Column A (Process)	Column B (Term)
(a) Mineralization	(1) Conversion of organic to inorganic nutrients
(b) Humification	(2) Formation of stable humus
(c) Photosynthesis	(3) Conversion of sunlight to chemical energy
(d) Respiration	(4) Energy release from food molecules

A) 1, (b) 2, (c) 3, (d) 4

B) 2, (b) 1, (c) 4, (d) 3

C) 3, (b) 2, (c) 1, (d) 4

D) 4, (b) 3, (c) 2, (d) 1

168.

Match the following types of organisms with their examples:

Column A (Organism)	Column B (Example)
(a) Producers	(1) Fungi
(b) Decomposers	(2) Algae
(c) Herbivores	(3) Grasshopper
(d) Carnivores	(4) Lion

A) 2, (b) 1, (c) 3, (d) 4

B) 3, (b) 4, (c) 2, (d) 1

C) 1, (b) 2, (c) 4, (d) 3

D) 4, (b) 3, (c) 1, (d) 2

169.

Match the following water bodies with their ecosystem types:

Column A (Water body)	Column B (Ecosystem type)
(a) Ocean	(1) Marine ecosystem
(b) Pond	(2) Freshwater ecosystem
(c) Estuary	(3) Brackish water ecosystem
(d) River	(4) Freshwater ecosystem

A) 1, (b) 2, (c) 3, (d) 4

B) 2, (b) 1, (c) 4, (d) 3

C) 3, (b) 4, (c) 1, (d) 2

D) 4, (b) 3, (c) 2, (d) 1

170.

Match the following terms related to ecosystem productivity:

Column A	Column B
(a) GPP (Gross primary productivity)	(1) Total energy produced by green plants
(b) NPP (Net primary productivity)	(2) Energy stored after plant respiration
(c) Secondary productivity	(4) Productivity of herbivores and carnivores
(d) Respiration	(3) Energy used by plants for metabolic activities

A) 1, (b) 2, (c) 4, (d) 3

B) 2, (b) 3, (c) 4, (d) 1

C) 3, (b) 4, (c) 1, (d) 2

D) 4, (b) 1, (c) 2, (d) 3

171.

Match the following types of consumers with their example:

Column A	Column B
(a) Primary consumer	(1) Bear
(b) Secondary consumer	(2) Snake
(c) Tertiary consumer	(3) Hawk
(d) Omnivore	(4) Deer

A) 1, (b) 2, (c) 3, (d) 4

B) 2, (b) 3, (c) 4, (d) 1

C) 3, (b) 4, (c) 1, (d) 2

D) 4, (b) 2, (c) 3, (d) 1

172.

Match the following ecosystem terms with their examples:

Column A	Column B
(a) Pond ecosystem	(1) Biotic and abiotic features
(b) Food web	(2) Interconnected food chains
(c) Niche	(3) Role of species in ecosystem
(d) Habitat	(4) Place where organism lives

A) 2, (b) 3, (c) 4, (d) 1

B) 3, (b) 4, (c) 1, (d) 2

C) 4, (b) 1, (c) 2, (d) 3

D) 1, (b) 2, (c) 3, (d) 4

173.

Choose the incorrect match

Column A	Column B
(a) Producers	(1) Convert solar energy to chemical energy
(b) Primary Consumers	(2) Feed on producers or other consumers
(c) Detritivores	(3) Convert living organisms into dead organic matter
(d) Decomposers	(4) Break down organic matter to inorganic form

A) a

B) b

C) c

D) d

174.

Pyramid drawn for food web is

A) upright pyramid of number

B) Upright pyramid of biomass

C) Inverted pyramid of energy

D) Not possible to draw

175.

Position of decomposers in food chain

A) Base

B) Next to producers

C) Top

D) No place

176.

Approximate percentage of energy transferred from one trophic level to the next in a food chain is:

A) 1%

B) 10%

C) 50%

D) 90%

177.

Which ecological pyramid is always upright?

A) Pyramid of numbers

B) Pyramid of biomass

C) Pyramid of energy

D) All three

178.

Net primary productivity (NPP) is equal to:

A) GPP + respiration by autotrophs

B) GPP – respiration by autotrophs

C) GPP – respiration by heterotrophs

D) GPP + respiration by heterotrophs

179.

Which food chain starts with dead organic matter and is especially important in ecosystems like forests and soils?

A) Grazing (herbivore) food chain

B) Detritus food chain

C) Parasitic food chain

D) Saprophytic food chain

180.

The usual unit for expressing primary productivity in ecosystems is:

A) kcal/m²/yr

B) gC/m²/yr

C) g/m³/yr

D) kg/ha/min

ANSWER KEY

Physics

1.	4	13.	2	25.	2	37.	3
2.	3	14.	1	26.	3	38.	3
3.	3	15.	1	27.	3	39.	2
4.	2	16.	2	28.	2	40.	4
5.	3	17.	1	29.	1	41.	2
6.	1	18.	4	30.	1	42.	1
7.	2	19.	3	31.	1	43.	1
8.	3	20.	1	32.	1	44.	2
9.	2	21.	4	33.	1	45.	1
10.	1	22.	1	34.	2		
11.	4	23.	3	35.	3		
12.	3	24.	2	36.	4		

Chemistry

46.	2	58.	2	70.	1	82.	1
47.	2	59.	2	71.	1	83.	1
48.	2	60.	3	72.	1	84.	4
49.	1	61.	2	73.	3	85.	1
50.	2	62.	1	74.	1	86.	3
51.	3	63.	4	75.	1	87.	4
52.	4	64.	2	76.	1	88.	1
53.	3	65.	1	77.	1	89.	1
54.	2	66.	3	78.	1	90.	1
55.	1	67.	1	79.	1		
56.	1	68.	1	80.	1		
57.	2	69.	1	81.	1		

Botany

91.	2	103.	1	115.	2	127.	1
92.	2	104.	2	116.	2	128.	2
93.	2	105.	3	117.	3	129.	3
94.	3	106.	2	118.	2	130.	2
95.	2	107.	3	119.	2	131.	2
96.	2	108.	4	120.	2	132.	1
97.	2	109.	1	121.	2	133.	2
98.	3	110.	3	122.	3	134.	1
99.	1	111.	2	123.	1	135.	2
100.	3	112.	1	124.	2		
101.	1	113.	1	125.	2		
102.	2	114.	2	126.	3		

Zoology

136.	1	148.	1	160.	1	172.	4
137.	2	149.	4	161.	3	173.	3
138.	1	150.	3	162.	1	174.	4
139.	4	151.	1	163.	1	175.	4
140.	2	152.	2	164.	1	176.	2
141.	1	153.	2	165.	1	177.	3
142.	1	154.	2	166.	1	178.	2
143.	1	155.	3	167.	1	179.	2
144.	2	156.	2	168.	1	180.	2
145.	2	157.	2	169.	1		
146.	1	158.	2	170.	1		
147.	3	159.	3	171.	4		

SOLUTIONS

Q1.

Answer: D

Explanation:

Assuming the blocks move together, the common acceleration is

$a = F/(m_A + m_B) = 80/20 = 4 \text{ m/s}^2$. The friction force required to push block B is $m_B a = 40 \text{ N}$, which is less than the maximum available friction $\mu m_A g = 50 \text{ N}$, so no slipping occurs and both blocks accelerate together at 4 m/s^2 .

Q2.

Answer: C

Explanation:

If the blocks moved together, the common acceleration would be

$a = F/(m_A + m_B) = 5 \text{ m/s}^2$, requiring a friction force of 50 N on block B. Since the maximum available friction is only $\mu m_A g = 40 \text{ N}$, the blocks slip. With kinetic friction $f = 40 \text{ N}$, block A accelerates at $a_A = (F - f)/m_A = 6 \text{ m/s}^2$ and block B at $a_B = f/m_B = 4 \text{ m/s}^2$.

Q3.

Answer: C

Explanation:

Assuming the blocks move together, the common acceleration is

$a = F/(m_A + m_B) = 120/30 = 4 \text{ m/s}^2$. The friction force required to drag block B is $m_B a = 40 \text{ N}$, which is well below the maximum available friction $\mu m_A g = 100 \text{ N}$, so no slipping occurs and both blocks accelerate together at 4 m/s^2 .

Q4.

Answer: B

Explanation:

The upper block A is driven only by friction, so its maximum possible acceleration is

$a_{max} = \mu g = 5 \text{ m/s}^2$. Sliding begins when the common acceleration $F/(m_A + m_B)$ just exceeds this value, giving $F_{min} = \mu g(m_A + m_B) = 0.5 \times 10 \times 40 = 200 \text{ N}$.

Q5.

Answer: C

Explanation:

Since the interface between A and B is frictionless ($\mu = 0$), no horizontal force is transmitted from A to B, so $f_{AB} = 0$. Block A therefore slides freely with

$a_A = F/m_A = 20/4 = 5 \text{ m/s}^2$, while block B, having no driving force acting on it, stays at rest and needs no friction from the ground, giving $a_B = 0$ and $f_{BG} = 0$.

Q6.

Answer: A

Explanation:

The maximum static friction between A and B is 30 N , which is less than the applied 42 N , so block A slides over B with kinetic friction $f_{AB} = 30 \text{ N}$, giving

$a_A = (42 - 30)/6 = 2 \text{ m/s}^2$. The reaction friction of 30 N pushing block B forward is far below the maximum static friction from the ground (90 N), so B remains at rest and the ground supplies an equal balancing static friction $f_{BG} = 30 \text{ N}$.

Q7.

Answer: B

Explanation:

The upper block A is driven only by friction, whose maximum value is $\mu m_A g = 30 \text{ N}$, limiting A to 6 m/s^2 . Testing the no-slip assumption gives a common acceleration of about 7.33 m/s^2 , which needs more friction than available, so A slips on B with $f_{AB} = 30 \text{ N}$ and $a_A = 6 \text{ m/s}^2$. Block B then has net force $170 - 30 - 60 = 80 \text{ N}$, giving $a_B = 8 \text{ m/s}^2$ with ground friction $f_{BG} = 60 \text{ N}$.

Q8.

Answer: C

Explanation:

The ground exerts kinetic friction $f_{BG} = \mu'(m_A + m_B)g = 60 \text{ N}$, so the net driving force on the system is $84 - 60 = 24 \text{ N}$, giving a common acceleration of $24/12 = 2 \text{ m/s}^2$ if the blocks move together. The friction needed to accelerate block A at this rate is only $m_A a = 8 \text{ N}$, well below the limiting value of 20 N , so no slipping occurs: both blocks move at 2 m/s^2 with $f_{AB} = 8 \text{ N}$ (static) and $f_{BG} = 60 \text{ N}$ (kinetic).

Q9.

Answer: B

Explanation:

The two blocks move together as a single system. First, calculate the friction between the lower block and the ground. Then determine the net force acting on the complete system.

Q10.

Answer: A

Explanation:

Since the ground is smooth, the only horizontal external force on the two-block system is 15 N . The blocks move together because the required friction is less than the maximum available friction. Hence, the acceleration of both blocks is 1.5 m/s^2 .

Q11.

Answer: D

Explanation:

The upper block tends to move to the right and exerts friction on the lower block towards the right. The maximum friction between the blocks is 16 N . This 16 N friction can be completely balanced by the static friction from the ground, whose maximum value is 30 N . Hence, the lower block remains at rest.

Q12.

Answer: C

Explanation:

The lower block moves to the right due to the applied force. The upper block is accelerated only by friction. Since the required friction for both blocks to move together is greater than the maximum friction available between the blocks, slipping occurs. Therefore, the friction on the upper block is maximum friction, giving its acceleration as 2 m/s^2 .

Q13.

Answer: B

Explanation:

Since the ground is smooth, the only external horizontal force on the two-block system is 45 N . The required friction between the blocks is less than the maximum possible friction, so both blocks move together with the same acceleration.

Q14.

Answer: A

Explanation:

First, determine if the system moves by comparing the applied force with the maximum friction from the ground. Then, assume the blocks move together to find a common acceleration and check if the required friction between the blocks exceeds the maximum static friction. Since it does, the blocks slip, and their accelerations must be calculated separately using kinetic friction.

Q15.

Answer: A

Explanation:

First, we determine the maximum limiting friction between the two blocks to see if they can move together. The maximum static friction is $f_{\max} = \mu m_1 g = 0.4 \times 4 \times 10 = 16\text{ N}$. Next, we assume the blocks move together as a single system. The total mass is $4 + 6 = 10\text{ kg}$. Since the ground is smooth, the only external force is the applied 20 N . The common acceleration would be $a = \frac{F}{m_{\text{total}}} = \frac{20}{10} = 2\text{ m/s}^2$. To maintain this acceleration for the lower block (6 kg), a force of $f_{\text{req}} = m_2 a = 6 \times 2 = 12\text{ N}$ is required. This force is provided by the friction between the blocks. Since the required friction (12 N) is less than the maximum possible friction (16 N), the blocks do not slip. They move together with an acceleration of 2 m/s^2 , and the frictional force is 12 N .

Q16.

Answer: B

Explanation:

Calculate the maximum static friction between the blocks and between the lower block and the ground. Compare the applied force and the transmitted frictional force to these limits to determine the motion of each block.

Q17.

Answer: A

Explanation:

Calculate the maximum static friction between the lower block and the ground. Compare this with the applied force to determine if the system moves. Since the applied force equals the maximum static friction, the system remains stationary.

Q18.

Answer: D

Explanation:

First, assume the blocks move together to find the common acceleration. Then, calculate the frictional force required to accelerate the lower block. Compare this required force with the maximum possible static friction to confirm if they move together.

Q19.

Answer: C

Explanation:

First, calculate the maximum frictional force between the lower block and the ground. The total normal force is $(2 + 3)g = 50$ N. The friction is $0.2 \times 50 = 10$ N. Since the applied force (30 N) is greater than this friction, the system moves. Next, assume both blocks move together. The net force is $30 - 10 = 20$ N. The total mass is 5 kg. The common acceleration is $20/5 = 4 \text{ m/s}^2$. Finally, check if the upper block can move with this acceleration. The force required is $m_1 a = 2 \times 4 = 8$ N. The maximum friction between the blocks is $\mu_1 m_1 g = 0.4 \times 20 = 8$ N. Since the required force equals the maximum friction, the blocks move together without slipping (limiting case). Thus, the acceleration of both blocks is 4 m/s^2 .

Q20.

Answer: A

Explanation:

First, calculate the maximum static frictional force between the two blocks. The normal force is the weight of the upper block ($3 \text{ kg} \times 10 \text{ m/s}^2 = 30$ N). The maximum friction is $f_{\max} = \mu N = 0.5 \times 30 = 15$ N. Next, assume both blocks move together as a single system. The total mass is $3 + 5 = 10$ kg? No, $3 + 5 = 8$ kg. The surface is smooth, so there is no external friction. The common acceleration would be $a = F/m_{\text{total}} = 18/8 = 2.25 \text{ m/s}^2$. To maintain this acceleration for the lower block (5 kg), a frictional force of $f = m_2 a = 5 \times 2.25 = 11.25$ N is required. Since the required friction (11.25 N) is less than the maximum possible friction (15 N), the blocks do not slip. They move together with an acceleration of 2.25 m/s^2 , and the frictional force is 11.25 N.

Q21.

Answer: D

Explanation:

First, calculate the maximum frictional forces. The limiting friction between the blocks is $f_1 = \mu_1 m_1 g = 0.5 \times 4 \times 10 = 20$ N. The limiting friction between the lower block and ground is $f_2 = \mu_2 (m_1 + m_2) g = 0.4 \times 10 \times 10 = 40$ N. Since the applied force (50 N) is greater than the friction between blocks (20 N), the upper block slides. The kinetic friction is 20 N. The acceleration of the upper block is $a_1 = (50 - 20)/4 = 7.5 \text{ m/s}^2$. The force driving the lower block is the friction from the upper block (20 N). Since this driving force (20 N) is less than the maximum friction from the ground (40 N), the lower block does not move ($a_2 = 0$).

Q22.

Answer: A

Explanation:

Since all surfaces are smooth (frictionless), there is no horizontal force transferred from the lower block to the upper block. Therefore, the upper block (1 kg) has no horizontal acceleration ($a = 0$). The lower block (4 kg) accelerates under the applied force of 20 N. The normal force between the blocks simply supports the weight of the upper block.

Q23.

Answer: C

Explanation:

First, calculate the maximum static frictional force between the lower block and the ground. The total normal force is the weight of both blocks: $N = (2 + 3) \times 10 = 50$ N. The maximum friction is $f_{\max} = \mu_2 N = 0.5 \times 50 = 25$ N. The applied horizontal force is $F = 16$ N. Since the applied force (16 N) is less than the maximum limiting friction from the ground (25 N), the system will not move. The static friction from the ground will adjust to 16 N to balance the applied force. Consequently, both blocks remain stationary, and their accelerations are zero.

Q24.

Answer: B

Explanation:

First, determine the maximum static frictional force between the two blocks. The normal force is the weight of the upper block ($2 \text{ kg} \times 10 \text{ m/s}^2 = 20 \text{ N}$). The maximum friction is $f_{\max} = \mu N = 0.5 \times 20 = 10 \text{ N}$. Next, check if the blocks move together. If they did, the common acceleration would be $a = F/(m_1 + m_2) = 20/5 = 4 \text{ m/s}^2$. The force required to accelerate the lower block (3 kg) at this rate is $F_{\text{req}} = 3 \times 4 = 12 \text{ N}$. Since the required force (12 N) exceeds the maximum friction (10 N), the blocks will slip relative to each other. The kinetic friction acting between the blocks is $f_k = 10 \text{ N}$. The acceleration of the upper block (2 kg) is $a_1 = (F - f_k)/m_1 = (20 - 10)/2 = 5 \text{ m/s}^2$. The acceleration of the lower block (3 kg) is $a_2 = f_k/m_2 = 10/3 \approx 3.33 \text{ m/s}^2$. Among the given options, 5 m/s^2 corresponds to the acceleration of the upper block.

Q25.

Answer: B

Explanation:

The problem asks for the acceleration of each block, implying they move together with a common acceleration. We treat the two blocks as a single system. The total mass is $3 \text{ kg} + 2 \text{ kg} = 5 \text{ kg}$. The ground is smooth, so there is no external friction. The applied force is 30 N. Using Newton's second law, $a = F/m_{\text{total}} = 30/5 = 6 \text{ m/s}^2$. (Note: Strictly speaking, one should check for slipping. Max friction $f = 0.4 \times 30 = 12 \text{ N}$. Force required to accelerate the 3 kg block at 6 m/s^2 is 18 N. Since $18 > 12$, the blocks would actually slip ($a_{\text{top}} = 4, a_{\text{bottom}} = 9$). However, since 4 and 9 are not options and 6 is, the question assumes common motion, likely implying the force is applied to the upper block where the limiting case holds perfectly: $30 - f = 3a, f = 2a \Rightarrow 30 = 5a \Rightarrow a = 6, f = 12$).

Q26.

Answer: C

Explanation:

To find the acceleration of the system, we consider the external forces acting on the combination of the two blocks. The internal friction between the blocks does not affect the motion of the center of mass of the system. The total mass of the system is $M = 1 \text{ kg} + 4 \text{ kg} = 5 \text{ kg}$. The normal force from the ground is $N = Mg = 5 \times 10 = 50 \text{ N}$. The frictional force between the lower block and the ground is $f = \mu_2 N = 0.2 \times 50 = 10 \text{ N}$. The net external force on the system is $F_{\text{net}} = F_{\text{applied}} - f = 50 - 10 = 40 \text{ N}$. The acceleration of the system (center of mass) is $a = F_{\text{net}}/M = 40/5 = 8 \text{ m/s}^2$.

Q27.

Answer: C

Explanation:

The problem asks for the acceleration of the upper block. In many such textbook problems, if the options suggest a common acceleration, we first check if the blocks move together. Assuming the blocks move together as a single system and ignoring the ground friction for the sake of matching the provided options (or assuming the applied force overcomes it in a simplified model): Total mass $M = 4 \text{ kg} + 6 \text{ kg} = 10 \text{ kg}$. Applied force $F = 40 \text{ N}$.

Common acceleration $a = F/M = 40/10 = 4 \text{ m/s}^2$. (Note: Strictly speaking, with the given coefficients, the maximum friction between blocks is $0.3 \times 40 = 12 \text{ N}$. The force required to accelerate the 4 kg block at 4 m/s^2 is not the right way to check. The force required to accelerate the 6 kg block at 4 m/s^2 is $6 \times 4 = 24 \text{ N}$, which exceeds the max friction 12 N, so they should slip. Also, ground friction is $0.4 \times 100 = 40 \text{ N}$, which balances the applied force. However, based on the options, the intended answer follows the simple system acceleration formula $a = F/m_{\text{total}}$.)

Q28.

Answer: B

Explanation:

Normal reactions: between the blocks $N_1 = 2g = 20 \text{ N}$; between the lower block and the ground $N_2 = (2 + 3)g = 50 \text{ N}$.

Limiting friction values:

$$f_{1,\max} = 0.4 \times 20 = 8 \text{ N} \quad (\text{between blocks}), \quad f_{2,\max} = 0.3 \times 50 = 15 \text{ N} \quad (\text{ground})$$

Test for common motion: if the blocks moved together, $a = \frac{45-15}{5} = 6 \text{ m/s}^2$, and the lower block would require $f_1 = 15 + 3(6) = 33 \text{ N} > 8 \text{ N}$, which is impossible. Hence the blocks slip, and the interface friction is kinetic: $f_1 = 8 \text{ N}$.

Individual accelerations:

$$\text{Upper block: } 45 - 8 = 2a_1 \implies a_1 = 18.5 \text{ m/s}^2$$

$$\text{Lower block: } f_1 = 8 \text{ N} < f_{2,\max} = 15 \text{ N} \implies a_2 = 0$$

Acceleration of the system (centre of mass):

$$a_{\text{system}} = \frac{m_1 a_1 + m_2 a_2}{m_1 + m_2} = \frac{2(18.5) + 3(0)}{5} = \frac{37}{5} = 7.4 \text{ m/s}^2$$

Hence the correct option is B.

Q29.

Answer: A

Explanation:

To find the maximum force F for which the blocks move together, we analyze the forces acting on each block. The blocks move together implies they have the same acceleration a . The limiting factor is the static friction between the two blocks. First, calculate the maximum static friction between the blocks ($f_{1,\max}$) and the maximum friction between the lower block and the ground ($f_{2,\max}$). $f_{1,\max} = \mu_1 m_1 g = 0.5 \times 2 \times 10 = 10$ N. $f_{2,\max} = \mu_2 (m_1 + m_2) g = 0.2 \times (2 + 3) \times 10 = 10$ N. The lower block (3 kg) is driven forward only by the friction force from the upper block (f_1). The opposing force is the ground friction (f_2). For the lower block to accelerate, f_1 must exceed f_2 . However, the maximum possible driving force $f_{1,\max}$ is 10 N, which is equal to the maximum resisting force $f_{2,\max}$ (10 N). Thus, the maximum acceleration of the lower block is $a = (f_{1,\max} - f_{2,\max})/m_2 = (10 - 10)/3 = 0$ m/s². Since the blocks move together, the upper block must also have acceleration $a = 0$. For the upper block: $F - f_1 = m_1 a \Rightarrow F - 10 = 2(0) \Rightarrow F = 10$ N. So, the maximum force is 10 N. For any $F > 10$ N, the upper block will slip over the lower block.

Q30.

Answer: A

Explanation:

First, we determine the maximum possible static frictional force between the two blocks. The normal force on the upper block is its weight, $N = 4 \text{ kg} \times 10 \text{ m/s}^2 = 40$ N. The maximum static friction is $f_{\max} = \mu N = 0.4 \times 40 = 16$ N. Next, we assume the blocks move together as a single system. The total mass is $4 \text{ kg} + 6 \text{ kg} = 10 \text{ kg}$. Since the ground is smooth, the only external horizontal force is the applied 40 N. The common acceleration is $a = F/m_{\text{total}} = 40 \text{ N}/10 \text{ kg} = 4 \text{ m/s}^2$. To maintain this acceleration for the upper block (4 kg), a net force of $F_{\text{req}} = ma = 4 \text{ kg} \times 4 \text{ m/s}^2 = 16$ N is required. This force is provided by the friction between the blocks. Since the required friction (16 N) is equal to the maximum limiting friction (16 N), the blocks move together without slipping. The frictional force is 16 N and the acceleration is 4 m/s^2 .

Q31.

Answer: A

Explanation:

To prevent slipping, both blocks must move with the same acceleration. The upper block (5 kg) is accelerated solely by the static friction force from the lower block. We first find the maximum acceleration the upper block can sustain without slipping. Then, we treat the two blocks as a single system to find the required applied force F , accounting for the friction from the ground.

Q32.

Answer: A

Explanation:

First, calculate the maximum static frictional forces. The limiting friction between the blocks is $f_1 = \mu_1 m_1 g = 0.6 \times 2 \times 10 = 12$ N. The limiting friction between the lower block and the ground is $f_2 = \mu_2 (m_1 + m_2) g = 0.3 \times 10 \times 10 = 30$ N. Since the applied force (30 N) is greater than the friction between the blocks (12 N), the upper block slides over the lower block. The kinetic friction is 12 N. The acceleration of the upper block is $a_1 = (30 - 12)/2 = 9 \text{ m/s}^2$. The force driving the lower block is the friction from the upper block (12 N). Since this driving force (12 N) is less than the maximum friction from the ground (30 N), the lower block does not move ($a_2 = 0$).

Q33.

Answer: A

Explanation:

First, calculate the maximum limiting friction between the lower block and the ground. The total normal force is the weight of both blocks: $N = (3 + 7) \times 10 = 100$ N. The maximum friction is $f_{\max} = \mu_2 N = 0.5 \times 100 = 50$ N. The applied horizontal force is $F = 16$ N. Since the applied force (16 N) is less than the maximum limiting friction from the ground (50 N), the system will not move. The static friction from the ground will adjust to 16 N to balance the applied force. Consequently, both blocks remain stationary, and their accelerations are zero. Since there is no acceleration and no external force on the upper block, the frictional force between the blocks is also zero.

Q34.

Answer: B

Explanation:

For the block to remain stopped, the upward friction force must balance the weight of the block. Since friction is $f = \mu N$ and the normal reaction equals the applied force ($N = F$), the condition $\mu F = mg$ gives $F = \frac{mg}{\mu} = \frac{3 \times 10}{0.6} = 50$ N.

Q35.

Answer: C

Explanation:

To just start the block moving, the applied force must overcome the limiting static friction. On a horizontal surface the normal reaction is $N = mg$, so the required force is $F = \mu_s N = \mu_s mg = 0.75 \times 8 \times 10 = 60$ N.

Q36.

Answer: D

Explanation:

Static friction is self-adjusting: it matches the applied force until the limiting value $f_{max} = \mu_s mg$ is reached. Here $f_{max} = 0.6 \times 4 \times 10 = 24$ N, which is greater than the applied force of 6 N, so the block stays at rest and the friction force simply equals the applied force, $f = 6$ N.

Q37.

Answer: C

Explanation:

Since the wall is smooth, it exerts only a horizontal normal force on the ladder and no vertical frictional force. Therefore vertical equilibrium of the ladder gives the floor normal reaction $N = W = 300$ N, and the maximum (limiting) friction at the floor is $f_{max} = \mu N = 0.4 \times 300 = 120$ N.

Q38.

Answer: C

Explanation:

The pressing force provides the normal reaction $N = 200$ N, so the limiting friction is $f_{max} = \mu N = 0.4 \times 200 = 80$ N. The weight of the body is only $mg = 3 \times 10 = 30$ N, which is less than f_{max} , so the body stays at rest and static friction simply balances the weight: $f = mg = 30$ N.

Q39.

Answer: B

Explanation:

Once the block slides, the opposing kinetic friction is $f = \mu mg = 0.3 \times 5 \times 10 = 15$ N. The net driving force is $F - f = 35 - 15 = 20$ N, so the acceleration is $a = \frac{F - \mu mg}{m} = \frac{20}{5} = 4$ m/s².

Q40.

Answer: D

Explanation:

Uniform velocity means the applied force exactly balances friction, so $f = \mu mg = 0.2 \times 3 \times 9.8 = 5.88$ N. The distance covered in 5 s is $d = vt = 2 \times 5 = 10$ m, so the work done against friction is $W = fd = 58.8$ J, and the heat generated is $H = \frac{W}{J} = \frac{58.8}{4.2} = 14$ cal.

Q41.

Answer: B

Explanation:

The applied force has horizontal and vertical components. The vertical component reduces the normal force, which in turn reduces the friction force. The net horizontal force is the horizontal component of the applied force minus the friction force. Using Newton's second law, we can find the acceleration.

Q42.

Answer: A

Explanation:

The net force acting on the body down the plane is the component of gravity ($mg \sin \theta$) minus the frictional force ($\mu mg \cos \theta$). Dividing this net force by the mass m gives the acceleration $a = g(\sin \theta - \mu \cos \theta)$.

Q43.

Answer: A

Explanation:

For the block to be just stationary, friction balances its weight: $f = W$ At limiting friction, $f = \mu N$ Since the applied horizontal force provides the normal reaction, $N = 16$ N Therefore, $W = \mu N = 0.25 \times 16 = 4$ N Hence, the correct answer is A) 4N.

Q44.

Answer: B

Explanation:

The 12kg block moves downward, the 8kg block moves upward, and the 10kg block moves toward the 12kg side. Net driving force: $F_{net} = (12 - 8)g = 4(10) = 40$ N Total mass: $M = 8 + 10 + 12 = 30$ kg Therefore, $a = \frac{40}{30} = 1.33$ m/s² Correction: Since the central 10kg block is connected to two separate strings, its acceleration is the same magnitude as the hanging blocks only when the string constraints are satisfied. Thus the correct equations must be solved individually. For the left block: $8g - T_1 = 8a$ For the right block: $12g - T_2 = 12a$ For the central block: $T_2 - T_1 = 10a$ Solving these equations gives: $a = 1.67$ m/s²

Q45.

Answer: A

Explanation:

For the 10kg block on the incline, $N = mg \cos 30^\circ$ $f = \mu N$ The hanging block pulls the inclined block upward. Therefore, friction acts down the incline. Using the equations of motion for both blocks: $100 - T = 10a$ $T - 50 - f = 10a$ Adding, $100 - 50 - f = 20a$ With $f = 0.2(10)(10) \cos 30^\circ = 17.3$ N we get $a = \frac{100 - 50 - 17.3}{20} = 1.63$ m/s² Hence, A.

Q46.

Answer: B

Explanation:

Calculating $\%C = \frac{12 \times 0.2 \times 100}{44 \times 0.3} = 18.18\%$ and $\%H = \frac{2 \times 0.1 \times 100}{18 \times 0.3} = 3.70\%$.

Q47.

Answer: B

Explanation:

Mass of HCl = 0.684 g. Since 36.5 g HCl reacts with 40 g NaOH, 0.684 g HCl reacts with 0.750 g (750 mg) NaOH. Unreacted NaOH = 1 g – 0.750 g = 250 mg.

Q48.

Answer: B

Explanation:

$\%C = 48\%$. Mole counts: $A = \frac{32}{64} = 0.5$, $B = \frac{20}{40} = 0.5$, $C = \frac{48}{32} = 1.5$. Simplest mole ratio is $1 : 1 : 3 \implies ABC_3$.

Q49.

Answer: A

Explanation:

Mass of pure $CaCO_3 = 20 \times \frac{20}{100} = 4$ g. Since 100 g $CaCO_3$ produces 44 g CO_2 , 4 g produces $\frac{44}{100} \times 4 = 1.76$ g CO_2 .

Q50.

Answer: B

Explanation:

Moles of HCl = $\frac{50}{1000} \times 0.5 = 0.025$ mol. Required pure $CaCO_3 = 0.0125$ mol = 1.25 g. Mass of 95% pure sample = $\frac{1.25}{0.95} \approx 1.32$ g.

Q51.

Answer: C

Explanation:

Moles of C = $\frac{78}{12} = 6.5$; Moles of H = $\frac{22}{1} = 22$. Mole ratio $C : H = 1 : 3.38 \approx 1 : 3 \implies CH_3$.

Q52.

Answer: D

Explanation:

Haber's reaction: $N_{2(g)} + 3H_{2(g)} \rightleftharpoons 2NH_{3(g)}$. Producing 20 moles NH_3 requires $\frac{3 \times 20}{2} = 30$ moles H_2 .

Q53.

Answer: C

Explanation:

$HCOOH$ yields 0.05 mol CO ; $H_2C_2O_4$ yields 0.05 mol $CO + 0.05$ mol CO_2 . KOH absorbs CO_2 . Remaining $CO = 0.10$ mol $\times 28 = 2.8$ g.

Q54.

Answer: B

Explanation:

8.45 g $AgNO_3$ (0.049 mol) reacts with 2.9 g $NaCl$ (0.049 mol) to yield 0.049 mol $AgCl$ precipitate = $0.049 \times 143.3 \approx 7$ g.

Q55.

Answer: A

Explanation:

Moles: $Mg = \frac{1.0}{24} = 0.0416$ mol; $O_2 = \frac{0.56}{32} = 0.0175$ mol. O_2 is limiting reagent. Reacted $Mg = 2 \times 0.0175 \times 24 = 0.84$ g. Excess $Mg = 1.0 - 0.84 = 0.16$ g.

Q56.

Answer: A

Explanation:

Moles: $H_2 = 1$ mol, $Cl_2 = 0.5$ mol (limiting reagent). 0.5 mol Cl_2 produces $2 \times 0.5 = 1$ mol $HCl_{(g)}$.

Q57.

Answer: B

Explanation:

Millimoles of chloride = 0.5; Millimoles of $AgNO_3 = 1.0$. Ratio is 2 : 1, so the chloride formula is XCl_2 .

Q58.

Answer: B

Explanation:

Molarity of $\text{Na}_2\text{CO}_3 = \frac{25.3 \times 1000}{106 \times 250} = 0.955 \text{ M}$. $[\text{Na}^+] = 2 \times 0.955 = 1.910 \text{ M}$ and $[\text{CO}_3^{2-}] = 0.955 \text{ M}$.

Q59.

Answer: B

Explanation:

Moles: $\text{H}_2 = 5 \text{ mol}$, $\text{O}_2 = 2 \text{ mol}$ (limiting reagent). 2 mol O_2 produces $4 \text{ mol H}_2\text{O}$.

Q60.

Answer: C

Explanation:

%O = 51.62%. Relative moles: C = 3.22, H = 9.67, O = 3.22. Ratio

C : H : O = 1 : 3 : 1 $\implies \text{CH}_3\text{O}$.

Q61.

Answer: B

Explanation:

Moles: $\text{PbO} = \frac{6.5}{224} = 0.029 \text{ mol}$ (limiting reagent). 0.029 mol PbO forms 0.029 mol PbCl_2 .

Q62.

Answer: A

Explanation:

Reaction: $\text{C}_3\text{H}_8 + 5\text{O}_2 \rightarrow 3\text{CO}_2 + 4\text{H}_2\text{O}$. $1 \text{ L of C}_3\text{H}_8$ requires 5 L of O_2 .

Q63.

Answer: D

Explanation:

Average mass = $\frac{200 \times 90 + 199 \times 8 + 202 \times 2}{100} = \frac{19996}{100} = 199.96 \text{ amu} \approx 200 \text{ amu}$.

Q64.

Answer: B

Explanation:

$\text{BaCO}_3 \rightarrow \text{BaO} + \text{CO}_2$. $9.85 \text{ g BaCO}_3 = 0.05 \text{ mol}$. Volume of $\text{CO}_2 = 0.05 \times 22.4 \text{ L} = 1.12 \text{ L}$ at NTP.

Q65.

Answer: A

Explanation:

1 mole NH_3 requires 1.25 moles O_2 . Since only 1 mole O_2 is available, O_2 is the limiting reagent and will be completely consumed.

Q66.

Answer: C

Explanation:

$\text{Zn} + \text{H}_2\text{SO}_4 \rightarrow \text{ZnSO}_4 + \text{H}_2 \uparrow$. 22400 mL H_2 is produced by 65 g Zn . 224 mL H_2 requires $\frac{65}{22400} \times 224 = 0.65 \text{ g Zn}$.

Q67.

Answer: A

Explanation:

From the balanced equation $\text{CH}_4 + 2\text{O}_2 \rightarrow \text{CO}_2 + 2\text{H}_2\text{O}$, 22 g of CO_2 corresponds to 0.5 moles . Therefore, $0.5 \text{ moles of CH}_4$ is required, which equals $0.5 \times 16 = 8 \text{ g}$.

Q68.

Answer: A

Explanation:

Moles of $\text{CO}_2 = \frac{22}{44} = 0.5 \text{ mol} = 50 \times 10^{-2} \text{ mol}$. Since 1 mole CH_4 produces 1 mole CO_2 , required moles of $\text{CH}_4 = 50 \times 10^{-2} \text{ mol}$, so $x = 50$.

Q69.

Answer: A

Explanation:

Let the weight of CaCO_3 be $x \text{ g}$. The residual oxides satisfy $\frac{56x}{100} + \frac{40(2.21-x)}{84} = 1.152 \text{ g}$, yielding $x \approx 1.187 \text{ g of CaCO}_3$ and $1.023 \text{ g of MgCO}_3$.

Q70.

Answer: A

Explanation:

PbCl_2 is the limiting reagent. According to the stoichiometry, amount of $\text{Pb}_3(\text{PO}_4)_2$ formed = $\frac{72}{3} = 24 \text{ mmol}$.

Q71.

Answer: A

Explanation:

Using $C_xH_y + \left(x + \frac{y}{4}\right)O_2 \rightarrow xCO_2 + \frac{y}{2}H_2O$, $10x = 40 \implies x = 4$, and $10 \times \frac{y}{2} = 50 \implies y = 10$. Total atoms = $4 + 10 = 14$.

Q72.

Answer: A

Explanation:

All four statements are correct based on n -factor = 0.79, mole counts $Fe^{2+} = 0.79$ and $Fe^{3+} = 0.14$, metal deficiency in ccp lattice, and 85%/15% composition.

Q73.

Answer: C

Explanation:

$W_1 = 0.25 \times 62 \times 0.5$ and $W_2 = 0.25 \times 62 \times 0.25$. Therefore, mass ratio $\frac{W_1}{W_2} = \frac{0.5}{0.25} = 2 : 1$.

Q74.

Answer: A

Explanation:

Relative moles $C : H = 7.15 : 14.2 = 1 : 2$ gives empirical formula CH_2 (mass 14). $n = \frac{84}{14} = 6 \implies C_6H_{12}$, so there are 12 hydrogen atoms per molecule.

Q75.

Answer: A

Explanation:

Using $C_xH_y + \left(x + \frac{y}{4}\right)O_2 \rightarrow xCO_2 + \frac{y}{2}H_2O$, $\frac{y}{2} = 3 \implies y = 6$, and $x + \frac{6}{4} = 9.5 \implies x = 8$. Formula is C_8H_6 .

Q76.

Answer: A

Explanation:

Mass of C = $\frac{12}{44} \times 0.792 = 0.216$ g. $\%C = \frac{0.216}{0.492} \times 100 = 43.9\% \approx 44\%$.

Q77.

Answer: A

Explanation:

Moles of $Zn = \frac{11.5}{65.4} = 0.1758$ mol = moles of H_2 . Volume = $0.1758 \times 22.7 = 3.99$ L ≈ 4 L.

Q78.

Answer: A

Explanation:

Moles $C = 1$, $O_2 = 1.5$. Carbon is limiting reagent, producing 1 mol $CO = 22.7$ L = 227×10^{-1} L.

Q79.

Answer: A

Explanation:

$3BaCl_2 + 2Na_3PO_4 \rightarrow Ba_3(PO_4)_2 + 6NaCl$. Na_3PO_4 is limiting reagent. 2 moles Na_3PO_4 yield 1 mole $Ba_3(PO_4)_2$.

Q80.

Answer: A

Explanation:

Mass of C = 0.3 g $\implies$ 0.025 mol C = moles CO_2 . Mass of $CO_2 = 0.025 \times 44 = 1.1$ g = 11×10^{-1} g, so $x = 11$.

Q81.

Answer: A

Explanation:

Total sugar = $50 + 200 = 250$ g in 700 g solution. Mass $\% = \frac{250}{700} \times 100 = 35.71\% \approx 36\%$.

Q82.

Answer: A

Explanation:

Mass of compound = 0.25 g. Mass of H = 0.014 g $\implies \%H = 5.6\% = 56 \times 10^{-1}\%$, so $x = 56$.

Q83.

Answer: A

Explanation:

Moles of M_2CO_3 = moles of CO_2 = 0.01 mol. Molar mass = $\frac{1\text{ g}}{0.01\text{ mol}} = 100\text{ g mol}^{-1}$.

Q84.

Answer: D

Q85.

Answer: A

Explanation:

Mass C = 0.2165 g (43.5%), mass H = 0.0492 g (9.9%). Mass O = 0.2263 g $\implies \%O = 46\%$.

Q86.

Answer: C

Explanation:

Fe_3O_4 is limiting reagent (20 mol). Moles of $Fe = 3 \times 20 = 60$ mol. Mass of $Fe = 60 \times 56 = 3360$ g.

Q87.

Answer: D

Explanation:

Mass C = 90 g $\implies 75\%$; mass H = 30 g $\implies 25\%$.

Q88.

Answer: A

Explanation:

Mass C = $\frac{12}{44} \times 0.20 = 0.0545$ g. $\%C = \frac{0.0545}{0.30} \times 100 = 18.18\% \approx 18\%$.

Q89.

Answer: A

Explanation:

Moles of $H_2O = \frac{81}{18} = 4.5$ mol $\implies$ moles $CH_4 = 2.25 = 225 \times 10^{-2}$ mol, so $x = 225$.

Q90.

Answer: A

Explanation:

Empirical formula is CH_2O . Matches CH_3COOH ($C_2H_4O_2$) and HCHO (CH_2O), so exactly 2 formulae agree.

Q91.

Answer: B

Q92.

Answer: B

Q93.

Answer: B

Q94.

Answer: C

Q95.

Answer: B

Q96.

Answer: B

Q97.

Answer: B

Q98.

Answer: C

Q99.

Answer: A

Q100.

Answer: C

Q101.

Answer: A

Q102.

Answer: B

Q103.

Answer: A

Q104.

Answer: B

Q105.

Answer: C

Q106.

Answer: B

Q107.

Answer: C

Q108.

Answer: D

Q109.

Answer: A

Q110.

Answer: C

Q111.

Answer: B

Q112.

Answer: A

Q113.

Answer: A

Q114.

Answer: B

Q115.

Answer: B

Q116.

Answer: B

Q117.

Answer: C

Q118.

Answer: B

Q119.

Answer: B

Q120.

Answer: B

Q121.

Answer: B

Q122.

Answer: C

Q123.

Answer: A

Q124.

Answer: B

Q125.

Answer: B

Q126.

Answer: C

Q127.

Answer: A

Q128.

Answer: B

Q129.

Answer: C

Q130.

Answer: B

Q131.

Answer: B

Q132.

Answer: A

Q133.

Answer: B

Q134.

Answer: A

Q135.

Answer: B

Q136.

Answer: A

Q137.

Answer: B

Q138.

Answer: A

Q139.

Answer: D

Q140.

Answer: B

Q141.

Answer: A

Q142.

Answer: A

Q143.

Answer: A

Q144.

Answer: B

Q145.

Answer: B

Q146.

Answer: A

Q147.

Answer: C

Q148.

Answer: A

Q149.

Answer: D

Q150.

Answer: C

Q151.

Answer: A

Q152.

Answer: B

Q153.

Answer: B

Q154.

Answer: B

Q155.

Answer: C

Q156.

Answer: B

Q157.

Answer: B

Q158.

Answer: B

Q159.

Answer: C

Q160.

Answer: A

Q161.

Answer: C

Q162.

Answer: A

Q163.

Answer: A

Q164.

Answer: A

Q165.

Answer: A

Q166.

Answer: A

Q167.

Answer: A

Q168.

Answer: A

Q169.

Answer: A

Q170.

Answer: A

Q171.

Answer: D

Q172.

Answer: D

Q173.

Answer: C

Q174.

Answer: D

Q175.

Answer: D

Q176.

Answer: B

Q177.

Answer: C

Q178.

Answer: B

Q179.

Answer: B

Q180.

Answer: B