

02

d and f Block Elements

General Characteristics :

Position in periodic Table

1. The d-block of the periodic table contains the elements of the groups 3-12 in which the d-orbitals are progressively filled in each of the four long periods.
2. There are four series of the d-block elements.

3d series

21 Sc Scandium	22 Ti Titanium	23 V Vanadium	24 Cr Chromium	25 Mn Manganese	26 Fe Iron	27 Co Cobalt	28 Ni Nickel	29 Cu Copper	30 Zn Zinc
-----------------------------	-----------------------------	----------------------------	-----------------------------	------------------------------	-------------------------	---------------------------	---------------------------	---------------------------	-------------------------

4d Series

39 Y Yttrium	40 Zr Zirconium	41 Nb Niobium	42 Mo Molybdenum	43 Tc Technetium	44 Ru Ruthenium	45 Rh Rhodium	46 Pd Palladium	47 Ag Silver	48 Cd Cadmium
---------------------------	------------------------------	----------------------------	-------------------------------	-------------------------------	------------------------------	----------------------------	------------------------------	---------------------------	----------------------------

5d Series

57 La Lanthanum	72 Hf Hafnium	73 Ta Tantalum	74 W Tungsten	75 Re Rhenium	76 Os Osmium	77 Ir Iridium	78 Pt Platinum	79 Au Gold	80 Hg Mercury
------------------------------	----------------------------	-----------------------------	----------------------------	----------------------------	---------------------------	----------------------------	-----------------------------	-------------------------	----------------------------

6d Series

89 Ac Actinium	104 Rf Rutherfordium	105 Db Dubnium	106 Sg Seaborgium	107 Bh Bohrium	108 Hs Hassium	109 Mt Meitnerium	110 Ds Darmstadtium	111 Rg Roentgenium	112 Cn Copernicium
-----------------------------	-----------------------------------	-----------------------------	--------------------------------	-----------------------------	-----------------------------	--------------------------------	----------------------------------	---------------------------------	---------------------------------

Q. Why d-block elements are called transition elements ?

- (i) Their ultimate & Penultimate shell is incomplete as well as they show variable oxidation state.
 - (ii) A transition element is defined as the one which has incompletely filled d-orbitals in its ground state or in any one of its oxidation state.
 - (iii) Zinc, Cadmium and Mercury of group 12 have full d^{10} configuration in their ground state as well as in their common oxidation states and hence, are not regarded as transition metals.
 - (iv) All transition elements are 'd' block elements, but all 'd' block elements are not transition-elements.
- All d-block elements are metals, they show all metallic properties**

Electronic Configuration of the d-block Elements :

(A) General Electronic Configurations

$$(n-1)d^{1-10} ns^{1-2}$$

The (n-1) stands for the inner d orbitals which may have one to ten electrons.

The (n) stands for the outer most s orbital which may have one or two electrons.

(B) Exceptions

(a) Above generalization has several exceptions because

- (i) The energy difference between (n-1)d and ns orbitals is very less.
- (ii) Half and completely filled sets of orbitals are relatively more stable.

(b) A consequence of this factor is reflected in the electronic configuration of following elements.

(i) In 3d series — Cr and Cu

(ii) In 4d series — Nb, Mo, Tc, Ru, Rh, Pd, and Ag

(iii) In 5d series — Pt and Au

(iv) In 6d series — Rg

(C) Electronic Configuration of all d-block Elements.

(a) 3d-series

Sc	→	$18[\text{Ar}]$	$3d^1$	$4s^2$
Ti	→	$18[\text{Ar}]$	$3d^2$	$4s^2$
V	→	$18[\text{Ar}]$	$3d^3$	$4s^2$
Cr	→	$18[\text{Ar}]$	$3d^5$	$4s^1$
Mn	→	$18[\text{Ar}]$	$3d^5$	$4s^2$
Fe	→	$18[\text{Ar}]$	$3d^6$	$4s^2$
Co	→	$18[\text{Ar}]$	$3d^7$	$4s^2$
Ni	→	$18[\text{Ar}]$	$3d^8$	$4s^2$
Cu	→	$18[\text{Ar}]$	$3d^{10}$	$4s^1$
Zn	→	$18[\text{Ar}]$	$3d^{10}$	$4s^2$

(b) 4d-series

Y	→	$36[\text{Kr}]$	$4d^1$	$5s^2$
Zr	→	$36[\text{Kr}]$	$4d^2$	$5s^2$
Nb	→	$36[\text{Kr}]$	$4d^4$	$5s^1$
Mo	→	$36[\text{Kr}]$	$4d^5$	$5s^1$
Tc	→	$36[\text{Kr}]$	$4d^6$	$5s^1$
Ru	→	$36[\text{Kr}]$	$4d^7$	$5s^1$
Rh	→	$36[\text{Kr}]$	$4d^8$	$5s^1$
Pd	→	$36[\text{Kr}]$	$4d^{10}$	$5s^0$
Ag	→	$36[\text{Kr}]$	$4d^{10}$	$5s^1$
Cd	→	$36[\text{Kr}]$	$4d^{10}$	$5s^2$

(c) 5d-series

La	→	$54[\text{Xe}]$	$4f^{14}$	$5d^1$	$6s^2$
Hf	→	$54[\text{Xe}]$	$4f^{14}$	$5d^2$	$6s^2$
Ta	→	$54[\text{Xe}]$	$4f^{14}$	$5d^3$	$6s^2$
W	→	$54[\text{Xe}]$	$4f^{14}$	$5d^4$	$6s^2$
Re	→	$54[\text{Xe}]$	$4f^{14}$	$5d^5$	$6s^2$
Os	→	$54[\text{Xe}]$	$4f^{14}$	$5d^6$	$6s^2$
Ir	→	$54[\text{Xe}]$	$4f^{14}$	$5d^7$	$6s^2$
Pt	→	$54[\text{Xe}]$	$4f^{14}$	$5d^9$	$6s^1$
Au	→	$54[\text{Xe}]$	$4f^{14}$	$5d^{10}$	$6s^1$
Hg	→	$54[\text{Xe}]$	$4f^{14}$	$5d^{10}$	$6s^2$

(d) 6d-Series

Ac	→	$86[\text{Rn}]$	$5f^{14}$	$6d^1$	$7s^2$
Rf	→	$86[\text{Rn}]$	$5f^{14}$	$6d^2$	$7s^2$
Db	→	$86[\text{Rn}]$	$5f^{14}$	$6d^3$	$7s^2$
Sg	→	$86[\text{Rn}]$	$5f^{14}$	$6d^4$	$7s^2$
Bh	→	$86[\text{Rn}]$	$5f^{14}$	$6d^5$	$7s^2$
Hs	→	$86[\text{Rn}]$	$5f^{14}$	$6d^6$	$7s^2$
Mt	→	$86[\text{Rn}]$	$5f^{14}$	$6d^7$	$7s^2$
Ds	→	$86[\text{Rn}]$	$5f^{14}$	$6d^8$	$7s^2$
Rg	→	$86[\text{Rn}]$	$5f^{14}$	$6d^{10}$	$7s^1$
Cn	→	$86[\text{Rn}]$	$5f^{14}$	$6d^{10}$	$7s^2$

Atomic Radius :

In period

In initial five elements attraction between nucleus and valence shell remains more as compared to repulsion between valence shell and penultimate shell that's why atomic radius decreases from Sc to Mn.


In next three elements attraction between nucleus and valence shell becomes equal to repulsion between valence shell and penultimate shell that's why atomic radius of Fe, Co and Ni remains almost equal.

In last two elements attraction between nucleus and valence shell remains less as compared to repulsion between valence shell and penultimate shell that's why atomic radius from Cu to Zn increases.

In group

Atomic radii of 4d series elements are more than 3d series elements.

4d and 5d series elements nearly same atomic radii due to lanthanide contraction.



Trends in atomic radii of transition elements

Magnetic Properties :

- (i) **Paramagnetic substances.** The substances which are attracted by magnetic field are called paramagnetic substances and this character arises due to the presence of unpaired electrons in the atomic orbitals.
- (ii) **Diamagnetic substances.** The substances which are repelled by magnetic field are called diamagnetic substances and this character arises due to the presence of paired electrons in the atomic orbitals.
- (iii) **Ferromagnetism.** Substances attracted very strongly are said to be ferromagnetic, extreme form of paramagnetic.

Ferromagnetic materials $\Rightarrow$ Fe, Co, Ni

Most of the transition metal ions or their compounds have unpaired electrons in d-sub-shell (from configuration d^1 to d^9) and therefore, they give rise to paramagnetic character.

Metals, which have unpaired electrons show para-magnetism.




Spin only magnetic moment $\mu = \sqrt{n(n+2)}$

here n = no. of unpaired electron

Unit : Bohr Magnetron (B.M.)

Formation of Interstitial Compounds :

- H, C, N trapped inside Crystal lattice of metals
- Non-stoichiometric
- TiC , Mn_4N , Fe_3H , $\text{VH}_{0.56}$, $\text{TiH}_{1.7}$
- Retain metallic conductivity
- High melting point than pure metal
- They are very hard
- They are chemically inert

**Formation of Alloys :**

- (i) Transition elements have maximum tendency to form alloys.
- (ii) The reactivity of transition elements is very less and their sizes are almost similar. Due to this a transition metal atom in the lattice can be easily replaced by other transition metal atom and hence they have maximum tendency to form alloys.
- (iii) In the alloys, ratio of component metals is fixed.
- (iv) These are extremely hard and have high melting point.



Some important alloys

(a)	Bronze	Cu + Sn
(b)	Brass	Cu + Zn
(c)	Gun metal	Cu + Zn + Sn
(d)	German Silver	Cu + Zn + Ni
(e)	Stainless Steel	Cr + Ni
(f)	Invar	Fe + Ni
(g)	Alnico	Al + Ni + Co
(h)	Duralumin	Cu + Al + Mn
(i)	22 Carrot gold	Au + Ag
(j)	18 Carrot gold	Au + Ag + Cu
(k)	Silver UK coins	Cu + Ni
(l)	White Metal	Li + Pb
(m)	Solder	Sn + Pb
(n)	Nichrome	Ni + Cr + Fe
(o)	Bell Metal	Cu + Sn

Amalgam is a semisolid alloy which is formed by mixing a metal with Hg. Fe, Co, Ni do not form amalgam due to large difference in size.

Formation of Complex Compounds :

- Small size
- High charge density
- Availability of vacant orbital

Example :**Melting Point :**

(i) Melting and boiling point of d-block elements is more than that of s block elements.

Reason : Stronger metallic bond formed by present unpaired d-electrons.

(ii) In Zn, Cd, and Hg there is no unpaired electron present in d-orbital, hence due to absence of covalent bond melting and boiling point are very low in series. (Volatile metals Zn, Cd, Hg)


(iii) In 3d series Sc to Cr melting and boiling point increases then Mn to Zn melting and boiling point decreases

(iv) As the number of unpaired d-electron increases, the number of covalent bond and bond energy between the atoms is expected to increase up to Cr-Mo-W family where each of the d-orbital has only unpaired electron and the opportunity for covalent sharing is greatest.

(v) Mn and Tc have comparatively low melting point, due to weak metallic bond because of stable Half filled (d^5) configuration and high IP.

(vi) Lowest melting point Hg (-38°C); Highest melting point W (3400°C)


Trends in melting points of transition elements



Heat of Atomisation :

- Enthalpy of atomization means the energy required to break the metallic bond.
- Stronger is the metallic bond more will be the heat of atomisation.
- More number of unpaired electrons will favour the formation of metallic bond.
- In case of Mn due to stable $(n-1)d^5 ns^2$ configuration metallic bond is weak.
- Mn and Zn has low heat of atomisation due to stable configuration electron are not available for metallic bonding.

Heat of Atomisation



Catalytic Properties :

- The transition metals and their compounds are known for their catalytic activity.
- This is due to -

Multiple oxidation states & presence of vacant orbital.

Catalyst	Used
Fe	Haber's process for manufacture of NH_3
V_2O_5	Contact process for H_2SO_4 manufacture
Pt	Ostwald's process of nitric acid
Ni	Hydrogenation of oils.

Density :

- (i) The atomic volumes of the transition elements are low compared with the elements of group 1 and 2. This is because the increased effective nuclear charge. Consequently the densities of the transition metals are high.
- (ii) Across a period from left to right atomic volumes decrease and atomic masses increase. Hence the densities increases across a period.

Sc < Ti < V < Cr < Mn < Fe < Co < Ni = Cu > Zn (Zn has lower density because of its large atomic volume)

Minimum density in 3d series → **Sc**

Maximum density in 3d series → **Ni** and **Cu**

In Group : **3d < 4d < 5d**

Elements with highest densities are

Osmium (Os) = 22.51 g/cm³, Iridium (Ir) = 22.61 g/cm³

Oxidation States :

1. d-block element shows variable oxidation state due to partially filled configuration.
2. It has two types of oxidation states -
 - (a) Minimum oxidation state = ns electron
 - (b) Maximum oxidation state = ns electron + (n-1)d unpaired electron
3. The variability of oxidation states, a characteristic of transition elements arises out of incomplete filling of d-orbitals in such a way that their oxidation states differ from each other by unity.

Example :

V⁺² V⁺³ V⁺⁴ V⁺⁵

4. In d-block elements the higher oxidation states are favoured by the heavier members.

Example :

In group - 6, Mo (VI) and W(VI) are found to be more stable than Cr (VI).

Thus Cr (VI) in form of dichromate (Cr₂O₇²⁻) in acidic medium is a strong oxidizing agent, whereas MoO₃ and WO₃ are not.

5. Low oxidation states are found when a complex compound has ligand capable of π-acceptor character in addition to the σ-bonding. (synergic ligand like CO)

Example :

In Ni(CO)₄ and Fe(CO)₅, the oxidation state of nickel and iron is zero.

They show variable oxidation state.

Sc	Ti	V	Cr	Mn	Fe	Co	Ni	Cu	Zn
								+1	
	+2	+2	+2	<u>+2</u>	<u>+2</u>	<u>+2</u>	<u>+2</u>	<u>+2</u>	<u>+2</u>
<u>+3</u>	+3	<u>+3</u>	<u>+3</u>	+3	+3	+3	+3		
	<u>+4</u>	+4	+4	<u>+4</u>	+4	+4	+4		
		<u>+5</u>	+5	+5					
			<u>+6</u>	+6	+6				
				<u>+7</u>					

- Underlined states are the most stable ones.

Electrode Potential :

In addition to ionisation enthalpy, the other factors such as enthalpy of sublimation, hydration enthalpy etc. determine the stability of a particular oxidation state in solution. This can be explained in terms of their electrode potential values. The oxidation potential of a metal involves the following process:




This process actually takes place in the following three steps as given in following flowchart :

The oxidation potential which gives the tendency of the overall change to occur, depends upon the net effect of these three steps. The overall energy change is

$$\Delta H = \Delta_{\text{sub}}H^\ominus + \Delta_{\text{IE}}H + \Delta_{\text{hyd}}H$$

If SOP is +ve that means oxidation will be easy.

If SRP is +ve that means reduction will be easy.

**Some important examples :**

1. $E_{M^{+2}/M}^\ominus \Rightarrow$ +ve only for Cu among 3d elements because hydration energy of Cu^{+2} is not enough to compensate for sublimation energy, IE_1 and IE_2 for Cu.
2. $E_{M^{+3}/M^{+2}}^\ominus$
 - (i) $E_{\text{Cr}^{+3}/\text{Cr}^{+2}}^\ominus = -\text{ve}$
that means Cr^{+2} acts as a reducing agent because Cr^{+3} has t_{2g} half filled stable configuration.
 - (ii) $E_{\text{Mn}^{+3}/\text{Mn}^{+2}}^\ominus = +\text{ve}$
that means Mn^{+3} acts as an oxidising agent due to stability of half filled (d^5) configuration.

Trends in Stability of Higher Oxidation States :

The highest oxidation states are shown generally among halides and oxides of transition elements.

(I) In halides of transition elements

- (i) The transition elements react with halogens at high temperatures to form transition metal halides.
- (ii) Since fluorine is the most electronegative element, the transition metals show highest oxidation states with fluorine. The highest oxidation states are found in TiX_4 (tetrahalides, $X = \text{F}, \text{Cl}, \text{Br}$ and I), VF_5 and CrF_6 .
- (iii) The +7 oxidation state for Mn is not shown by simple halides. However, MnO_3F is known in which the oxidation state of Mn is +7.
- (iv) The tendency of fluorine to stabilise the highest oxidation state is due to either higher lattice enthalpy as in case of CoF_3 or higher bond enthalpy terms for the higher covalent compounds. eg. VF_5 and CrF_6 .
- (v) Fluorides are relatively unstable in their low oxidation states. For example, vanadium form only VX_2 ($X = \text{Cl}, \text{Br}$ or I) and copper can form CuX ($X = \text{Cl}, \text{Br}, \text{I}$). On the other hand all copper (II) halides are known except the iodide. This is because, Cu^{2+} oxidises I^- to I_2 .

$$2\text{Cu}^{2+} + 4\text{I}^- \longrightarrow \text{Cu}_2\text{I}_2(\text{s}) + \text{I}_2$$
- (vi) It has been observed that many copper (I) compounds are unstable in aqueous solution and they undergo disproportionation.

$$2\text{Cu}^+ \longrightarrow \text{Cu}^{2+} + \text{Cu}$$

Copper in +2 oxidation state is more stable than in +1 oxidation state. This can be explained on the basis of much larger negative hydration enthalpy ($\Delta_{\text{hyd}}H$) of $\text{Cu}^{2+}(\text{aq})$ than Cu^+ , which is sufficiently high to compensate second ionisation enthalpy of copper.

(II) In metal oxides and oxocations.

- (i) The ability of oxygen to stabilize the highest oxidation state is demonstrated in their oxides. The highest oxidation states in their oxides coincides with the group number. For example, the highest oxidation state of scandium of group 3 is +3 in its oxides, Sc_2O_3 whereas the highest oxidation state of manganese of group 7 is +7, in Mn_2O_7 .
- (ii) Besides the oxides, oxocation of the metals also stabilise higher oxidation states. For example, V^{+5} as VO_2^+ , V^{+4} as VO^{2+} and Ti^{+4} as TiO^{2+} .
- (iii) It may be noted that the ability of oxygen to stabilise these high oxidation states exceeds that of fluorine. For example, manganese forms highest fluoride as MnF_4 whereas the highest oxide is Mn_2O_7 . This is due to the fact that oxygen has ability to form multiple bonds to metals.
- (iv) The transition elements in the +2 and +3 oxidation states mostly form ionic bonds whereas with higher oxidation states, the bonds are essentially covalent e.g. in MnO_4^- all bonds are covalent. In these higher oxides the acidic character is predominant. Thus CrO_3 gives H_2CrO_4 and $\text{H}_2\text{Cr}_2\text{O}_7$ and Mn_2O_7 gives HMnO_4 . **V_2O_5 is, however amphoteric though mainly acidic and with alkalis as well as acids gives VO_4^{3-} and VO_2^+ respectively.**

Element (M)	$E^\circ_{\text{M}^{3+}/\text{M}^{2+}}$	$E^\circ_{\text{M}^{2+}/\text{M}}$	Comments
Ti	-0.37	-1.63	+3 Oxidation state is more stable
V	-0.26	-1.18	+3 Oxidation state is more stable
Cr	-0.41	-0.9	+3 Oxidation state is more stable
Mn	1.57	-1.18	+2 Oxidation state is more stable
Fe	0.77	-0.44	+2 Oxidation state is more stable
Co	1.97	-0.28	+2 Oxidation state is more stable
Ni	-	-0.25	+3 Oxidation state does not exist for Ni
Cu	-	0.34	+3 Oxidation state does not exist for Cu. 0 Oxidation state is more stable
Zn	-	-0.76	+3 Oxidation state does not exist for Zn



BEGINNER'S BOX-1

1. Transition elements exhibit variable oxidation states in their compounds due to

- (1) Strong shielding of (n-1) d-electrons
- (2) Very small energy difference between (n-1)d and ns orbitals
- (3) Presence of unpaired electrons
- (4) High energy difference between (n-1)d and ns orbitals

CDF104

2. Which element do not show more than one oxidation state

- (1) Mn
- (2) Cr
- (3) Fe
- (4) Sc

CDF105

3. Consider the ions Fe^{2+} , Fe^{3+} , Cr^{2+} , Cr^{3+} , Mn^{2+} , Mn^{3+}

- (i) Strongest oxidising ion is

- (1) Fe^{2+}
- (2) Cr^{3+}
- (3) Mn^{3+}
- (4) Mn^{2+}

- (ii) Strongest reducing ion is


- (1) Fe^{+3}
- (2) Mn^{2+}
- (3) Cr^{2+}
- (4) Mn^{3+}

CDF106

4.	Which compound does exist (1) MnO_3F (2) $\text{K}_2[\text{CuI}_4]$ (3) MnF_7 (4) CrF_7	CDF107
5.	In the form of dichromate, Cr(VI) is a strong oxidising agent in acidic medium but Mo(VI) in MoO_3 and W(VI) in WO_3 are not because (1) Cr(VI) is more stable than Mo(VI) & W(VI) (2) Mo(VI) and W(VI) are less stable than Cr(VI) (3) H.O.S. of heavier members of group 6 of transition metals are more stable (4) L.O.S. of heavier members of group 6 of transition metals are more stable	CDF108
6.	Which of the following show +8 oxidation state. (1) Fe, Ru (2) Fe, Os (3) Ru, Os (4) Mo, W	CDF109
7.	Which catalyst is used in contact process ? (1) Ni (2) V_2O_5 (3) Fe (4) $\text{AlCl}_3 + \text{TiCl}_4$	CDF110
8.	The element with maximum and minimum melting points in second transition series respectively. (1) Cr and Zn (2) Cr and Hg (3) Cr and Cd (4) Mo and Cd	CDF111
9.	Which of the following metal ions is expected to be paramagnetic? (1) Zn^{2+} (2) Ti^{3+} (3) Sc^{3+} (4) Ti^{4+}	CDF112
10.	Which of the element have highest density – (1) W (2) Os (3) Tc (4) Mo	CDF113

Potassium Dichromate $\text{K}_2\text{Cr}_2\text{O}_7$:**Preparation :**


- Dichromates are generally prepared from chromate, which in turn are obtained by the fusion of chromite ore (FeCr_2O_4) with sodium or potassium carbonate in free access of air.
- The reaction with sodium carbonate occurs as follows:



The yellow solution of sodium chromate is filtered and acidified with sulphuric acid to give a solution from which orange sodium dichromate, $\text{Na}_2\text{Cr}_2\text{O}_7 \cdot 2\text{H}_2\text{O}$ can be crystallised.



- Sodium dichromate is more soluble than potassium dichromate. The latter is therefore, prepared by treating the solution of sodium dichromate with potassium chloride.




Orange crystals of potassium dichromate crystallise out. The chromates and dichromates are interconvertible in aqueous solution depending upon pH of the solution. The oxidation state of chromium in chromate and dichromate is the same.




- $\text{K}_2\text{Cr}_2\text{O}_7$ is preferred over $\text{Na}_2\text{Cr}_2\text{O}_7$ as a primary standard in volumetric analysis (titration) because $\text{Na}_2\text{Cr}_2\text{O}_7$ is hygroscopic in nature but $\text{K}_2\text{Cr}_2\text{O}_7$ is not.

- The structures of chromate ion, CrO_4^{2-} and the dichromate ion, $\text{Cr}_2\text{O}_7^{2-}$ are shown below. The chromate ion is tetrahedral whereas the dichromate ion consists of two tetrahedra, sharing one corner with Cr-O-Cr bond angle of 126° .





Oxidising Property :-

Acidified $\text{K}_2\text{Cr}_2\text{O}_7$ (Orange)



Chromyl chloride test (used to identify ionic chlorides)

Generally Ionic or water soluble chloride gives chromyl chloride test.




AgCl , HgCl_2 & PbCl_2 are covalent or water insoluble chloride so they can't give chromyl chloride test.


Potassium Permanganate (KMnO_4) :

Preparation :


- Potassium permanganate is prepared by fusion of MnO_2 (Pyrolusite) with an alkali metal hydroxide and an oxidising agent like KNO_3 . This produces the dark green K_2MnO_4 which disproportionate in a neutral or acidic solution to give permanganate.



- Commercially it is prepared by the alkaline oxidative fusion of MnO_2 followed by the electrolytic oxidation of manganate (VI).



- In the laboratory, a manganese (II) ion salt is oxidised by peroxodisulphate to permanganate.



Physical Property :




- Potassium permanganate forms **dark purple (almost black)** crystals which are isostructural with those of KClO_4 . The salt is not very soluble in water (6.4 g/100 g of water at 293 K), but when heated it decomposes at 513 K.



The manganate and permanganate ions are tetrahedral; the π bonding takes place by overlap of p orbitals of oxygen with d orbitals of manganese.

The green manganate is paramagnetic because of one unpaired electron but the permanganate is diamagnetic due to the absence of unpaired electron.

Oxidising Property :



f-BLOCK ELEMENTS**Introduction :**

- (a) The elements constituting the f – block are those in which the 4f and 5f orbitals are progressively filled.
- (b) These elements are formal members of group 3 from which they have been taken out to form a separate f-block of the periodic table.
- (c) The name inner transition metals are often used to refer to the elements of f-block.
- (d) The f-block consists of the two series

(i) Lanthanides

The fourteen elements following lanthanum.

Because lanthanum closely resembles the lanthanides, it is usually included in any discussion of the lanthanides.

(ii) Actinides

The fourteen elements following actinium.

Because actinium closely resembles the actinides, it is usually included in any discussion of the actinides.


The lanthanides :

Name	Electronic Configuration
$^{58}\text{Ce} \longrightarrow$	$^{54}[\text{Xe}] 4f^1 5d^1 6s^2$ (Cerium)
$^{59}\text{Pr} \longrightarrow$	$^{54}[\text{Xe}] 4f^3 5d^0 6s^2$ (Praseodymium)
$^{60}\text{Nd} \longrightarrow$	$^{54}[\text{Xe}] 4f^4 5d^0 6s^2$ (Neodymium)
$^{61}\text{Pm} \longrightarrow$	$^{54}[\text{Xe}] 4f^5 5d^0 6s^2$ (Promethium)
$^{62}\text{Sm} \longrightarrow$	$^{54}[\text{Xe}] 4f^6 5d^0 6s^2$ (Samarium)
$^{63}\text{Eu} \longrightarrow$	$^{54}[\text{Xe}] 4f^7 5d^0 6s^2$ (Europium)
$^{64}\text{Gd} \longrightarrow$	$^{54}[\text{Xe}] 4f^7 5d^1 6s^2$ (Gadolinium)
$^{65}\text{Tb} \longrightarrow$	$^{54}[\text{Xe}] 4f^9 5d^0 6s^2$ (Terbium)
$^{66}\text{Dy} \longrightarrow$	$^{54}[\text{Xe}] 4f^{10} 5d^0 6s^2$ (Dysprosium)
$^{67}\text{Ho} \longrightarrow$	$^{54}[\text{Xe}] 4f^{11} 5d^0 6s^2$ (Holmium)
$^{68}\text{Er} \longrightarrow$	$^{54}[\text{Xe}] 4f^{12} 5d^0 6s^2$ (Erbium)
$^{69}\text{Tm} \longrightarrow$	$^{54}[\text{Xe}] 4f^{13} 5d^0 6s^2$ (Thulium)
$^{70}\text{Yb} \longrightarrow$	$^{54}[\text{Xe}] 4f^{14} 5d^0 6s^2$ (Ytterbium)
$^{71}\text{Lu} \longrightarrow$	$^{54}[\text{Xe}] 4f^{14} 5d^1 6s^2$ (Lutetium)

- It may be noted that atoms of these elements have electronic configuration with $6s^2$ common but variable occupancy of 4f level.
- The irregularities in the electronic configuration of lanthanides are related to the stabilities of the f^0 , f^7 , f^{14} occupancies of the 4f orbitals.

Atomic and Ionic Radius :

1. The overall decrease in ionic radii from lanthanum to lutetium is a unique feature in the chemistry of the lanthanides.
2. This is due to poor shielding of nucleus by inner penultimate 4f orbital.
3. The cumulative effect of the contraction in size of the lanthanide series, known as lanthanide contraction.
4. The decrease in atomic radii is not quite regular.
5. The decrease in ionic radii of M^{3+} comparatively more regular.



Trends in ionic radii of lanthanides

Consequences of lanthanide Contraction :

- Small size difference between size of 4d and 5d series of transition element.
- Effect of basic strength, Basic strength decreases on moving left to right.
- Steady decrease in size.


Oxidation States :

1. The common oxidation state for lanthanides is +3 but it also shows +2 and +4 state.
2. (i) Ce $\rightarrow$ +4 and +3. Ce⁺⁴ act as a oxidant & used in volumetric Analysis.
 (ii) Pr, Nd, Tb and Dy also exhibit +4 state but only in oxides, MO₂.
 (iii) Eu²⁺ is a strong reducing agent changing to the common +3 state.
 (iv) Yb²⁺ has f¹⁴ configuration is a reductant.
 (v) Tb⁺⁴ has f⁷ configuration is a oxidant.
 (vi) The behaviour of samarium is very much like europium, exhibiting both +2 and +3 oxidation states.

General Characteristics :

1. Silvery white
2. Soft metals
3. Tarnish rapidly in air
4. Typical metallic structure
5. Good conductors of heat
6. Good conductors of electricity
7. (i) Many trivalent lanthanide ions are coloured both in the solid state and in aqueous solutions.
 (ii) Colour of these ions may be attributed to the presence of unpaired f electrons. However, absorption bands are narrow, probably because of the excitation within f orbitals, called f-f transition.
 (iii) La³⁺ and Lu³⁺ ions does not show any colour but the rest do so.

8. (i) The lanthanide ions other than the f^0 type (La^{3+} and Ce^{4+}) and the f^{14} type (Yb^{2+} and Lu^{3+}) are all paramagnetic.
- (ii) The paramagnetism rises to maximum in Nd.



Other Properties :

Alloy Formation

Alloy of lanthanide with iron called misch metal (composition) $\rightarrow$ Ce – 50%, La – 25%, Nd – 15% & 10% other rare – earth metal & iron or Lanthanoid metal (95%) + Iron (5%)

The Actinides :

Name	Electronic configuration
${}_{90}\text{Th} \longrightarrow$	${}_{86}[\text{Rn}] 5f^0 6d^2 7s^2$ (Thorium)
${}_{91}\text{Pa} \longrightarrow$	${}_{86}[\text{Rn}] 5f^2 6d^1 7s^2$ (Protactinium)
${}_{92}\text{U} \longrightarrow$	${}_{86}[\text{Rn}] 5f^3 6d^1 7s^2$ (Uranium)
${}_{93}\text{Np} \longrightarrow$	${}_{86}[\text{Rn}] 5f^4 6d^1 7s^2$ (Neptunium)
${}_{94}\text{Pu} \longrightarrow$	${}_{86}[\text{Rn}] 5f^6 6d^0 7s^2$ (Plutonium)
${}_{95}\text{Am} \longrightarrow$	${}_{86}[\text{Rn}] 5f^7 6d^0 7s^2$ (Americium)
${}_{96}\text{Cm} \longrightarrow$	${}_{86}[\text{Rn}] 5f^7 6d^1 7s^2$ (Curium)
${}_{97}\text{Bk} \longrightarrow$	${}_{86}[\text{Rn}] 5f^9 6d^0 7s^2$ (Berkelium)
${}_{98}\text{Cf} \longrightarrow$	${}_{86}[\text{Rn}] 5f^{10} 6d^0 7s^2$ (Californium)
${}_{99}\text{Es} \longrightarrow$	${}_{86}[\text{Rn}] 5f^{11} 6d^0 7s^2$ (Einsteinium)
${}_{100}\text{Fm} \longrightarrow$	${}_{86}[\text{Rn}] 5f^{12} 6d^0 7s^2$ (Fermium)
${}_{101}\text{Md} \longrightarrow$	${}_{86}[\text{Rn}] 5f^{13} 6d^0 7s^2$ (Mendelevium)
${}_{102}\text{No} \longrightarrow$	${}_{86}[\text{Rn}] 5f^{14} 6d^0 7s^2$ (Nobelium)
${}_{103}\text{Lr} \longrightarrow$	${}_{86}[\text{Rn}] 5f^{14} 6d^1 7s^2$ (Lawrencium)

- It may be noted that atoms of these elements have electronic configuration with $7s^2$ common but variable occupancy of $5f$ and $6d$ sub shell.
- The irregularities in the electronic configuration of actinides are related to the stabilities of the f^0 , f^7 , f^{14} occupancies of the $5f$ orbitals.

Ionic Sizes :

- The general trend in lanthanides is observable in the actinides as well. There is a gradual decrease in the size of atoms or M^{3+} ions across the series. This may be referred to as the actinide contraction (like lanthanide contraction). The contraction is, however, greater from element to element in this series resulting from poor shielding by 5f electrons.

Actinide Contraction :

- Due to poor shielding of 5f orbitals. Actinide contraction is more effective than lanthanide contraction.

Colour :

- Actinides are generally coloured.
- The colour of actinides depends upon the number of 5f electrons.
- If there is no unpaired electrons it will be colourless.

Oxidation State :

⇒ Most common oxidation state is +3 beside that they show +4, +5, +6 oxidation state in certain elements.

Oxidation State of Lanthanoids

Ce	Pr	Nd	Pm	Sm	Eu	Gd	Tb	Dy	Ho	Er	Tm	Yb	Lu
				+2	+2						+2	+2	
+3	+3	+3	+3	+3	+3	+3	+3	+3	+3	+3	+3	+3	+3
+4	+4						+4						

Oxidation State of Actinoids

Th	Pa	U	Np	Pu	Am	Cm	Bk	Cf	Es	Fm	Md	No	Lr
+3	+3	+3	+3	+3	+3	+3	+3	+3	+3	+3	+3	+3	+3
+4	+4	+4	+4	+4	+4	+4	+4						
	+5	+5	+5	+5	+5								
		+6	+6	+6	+6								
			+7	+7									

Complex Formation :

- Actinides have stronger tendency towards complex formation.

Radioactivity :

- All actinides are radioactive while lanthanides are non-radioactive [except promethium (pm)].
- Oxides and hydroxides of actinides are more basic than lanthanide.



BEGINNER'S BOX-2

- $X \xrightarrow[\text{K}_2\text{Cr}_2\text{O}_7]{\text{conc. H}_2\text{SO}_4} \text{red brown vapours. X can be}$

(1) NaI (2) NaBr (3) Both (1) and (2) (4) None of these

CDF114
- $\text{X} \xrightarrow[\text{air}]{\text{KOH}} \text{Y} \xrightarrow[\text{an oxidising agent}]{\text{H}^+} \text{Z} + \text{X}$

black coloured ore dark green an oxidising agent

True statement is

(1) X can't be MnO_2 (2) Y can be KMnO_4

(3) Z can be K_2MnO_4 (4) X is MnO_2 and Y is K_2MnO_4

CDF115
- $\text{X} \xrightleftharpoons[\text{OH}^-]{\text{H}^+} \text{Y}; \text{Y} \xrightarrow{\text{SO}_2} \text{Z} + \text{SO}_4^{2-}$

X and Y are oxy anion of Cr than oxidation state of Cr in X, Y, Z will be respectively

(1) +3, +6, +6 (2) +6, +3, +6 (3) +6, +6, +6 (4) +6, +6, +3

CDF116
- When KMnO_4 solution is added to oxalic acid solution, the decolourisation is slow in beginning but become instantaneous after some time because

(1) CO_2 is a by product (2) Mn^{2+} acts as a catalyst

(3) Reaction is exothermic (4) MnO_4^- catalyzes the reaction

CDF117
- The elements from cerium (At. No. 58) to lutetium (At. No. 71) in which 4f energy levels are filled up are known as :

(1) Lanthanides (2) Rare earth metals (3) Lanthanones (4) All the above

CDF118
- Which of the following is radioactive element :

(1) Ce (2) Lu (3) Eu (4) Pm

CDF119
- Which of the following ion is paramagnetic ?

(1) Ce^{+4} (2) Lu^{+3} (3) Yb^{+2} (4) Eu^{+2}

CDF120
- Which of the following ion do not show colour?

(1) Lu^{3+} (2) Pm^{3+} (3) Eu^{3+} (4) Dy^{3+}

CDF121
- Which type of hydroxide generally formed by Lanthanoids ?

(1) $\text{Ln}(\text{OH})_3$ (2) $\text{Ln}(\text{OH})_2$ (3) $\text{Ln}(\text{OH})_4$ (4) $\text{Ln}(\text{OH})_6$

CDF122
- Highest oxidation state +7 shown by which actinoids ?

(1) Np, Pu (2) Pa, U (3) Am, Cm (4) Cf, Es

CDF123

NCERT QUESTIONS

Q.1 Why do the transition elements have higher boiling & melting points ?

Ans. Because of having larger number of unpaired electrons in their atoms, they have stronger interatomic interaction. Hence strong metallic bonds between the atoms of these elements attribute to their high melting and boiling points.

Q.2 Give the main difference in oxidation states of transition elements & p-block elements.


Ans. (i) The variation of oxidation state in transition elements is in such a way that their oxidation state differ from each other by unity e.g. V^{+2} , V^{+3} , V^{+4} , V^{+5} . While in p block elements oxidation states normally differ by two due to inert pair effect.

(ii) Stability of higher oxidation state in transition elements increase down the group
e.g. Pt^{+4} is more stable than Pt^{+2}

While in p block elements lower oxidation state become increasingly stable down the group as result of inert pair effect. e.g. Pb^{+2} is more stable than Pb^{+4} .

Q.3 Explain why Cu^+ ion is not stable in aqueous solutions.

Ans. Cu^+ ions in aqueous solution undergoes disproportionation.



This is because that hydration energy of $Cu_{(aq)}^{+2}$ ions is sufficiently high to compensate IE_2 of Cu.

Q.4 Which is a stronger reducing agent Cr^{+2} or Fe^{+2} and why ?

Ans. Cr^{+2} is stronger reducing agent than Fe^{+2}



Reason : $d^4 \longrightarrow d^3$ occurs in case of $Cr^{+2} \longrightarrow Cr^{+3}$ but $d^6 \longrightarrow d^5$ occurs in case of $Fe^{+2} \longrightarrow Fe^{+3}$ in a medium like water d^3 is more stable as compared to d^5 (on the basis of CFSE value)

Q.5 In first transition series (3d series) the enthalpy of atomisation of zinc is lowest, why ?

Ans. In the formation of metallic bonds no e^- from 3d subshell is involved in case of Zn because all 3d orbitals are fulfilled. Which decrease strength of metallic bond as well as enthalpy of atomisation of Zn.

Q.6 How iron (III) catalyses the reaction between iodide & persulphate?

Ans. $2Fe^{3+} + 2I^- \longrightarrow 2Fe^{2+} + I_2$



BEGINNER'S BOX

ANSWERS KEY

BEGINNER'S BOX-1	Que.	1	2	3	4	5	6	7	8	9	10
	Ans.	2	4	3, 3	1	3	3	2	4	2	2

BEGINNER'S BOX-2	Que.	1	2	3	4	5	6	7	8	9	10
	Ans.	2	4	4	2	4	4	4	1	1	1