


05

Thermodynamics

Introduction :


Thermo (Heat/energy) + Dynamics (Flow/motion)

Thermodynamics is the branch of science which deals with the energy changes taking place in all physical and chemical processes but **Chemical thermodynamics** is the branch of thermodynamics which deals with the study of energy changes taking place in chemical processes.



Thermo is derived from a Greek word "Thermes" means Heat

Dynamics is derived from a Greek word "Dynamikos" means study of motion.



Limitations of thermodynamics :


- (i) Thermodynamics deals with the properties like temperature, pressure, volume, etc. of matter in bulk but doesn't tell us anything about the individual properties of atoms or molecules.

or


Thermodynamics deals with **macroscopic system** but not with **microscopic system**.

- (ii) It tells us whether a given chemical reaction will take place or not under the given set of conditions but doesn't tell us anything about the **rate of reaction**.

Some Important Terms :




-  **System :** A system is defined as a specific part of universe or specified portion of the matter which is under experimental investigation
-  **Surrounding :** The rest part of the universe excluding the system is called surrounding.
Universe = System + Surrounding
-  **Boundary :** Anything which separates system and surrounding is called boundary.



For example :

A reaction is carried out in a beaker. The contents of beaker constitute the system, beaker serves as boundary and anything which is outside the beaker is called surrounding.

Types of System :



Open system : This type of system can exchange energy as well as matter with the surrounding. The boundary is neither sealed nor insulated. Total mass will not remain constant.

Eg.1 Coffee in open glass, Hot water in open glass.

Eg.2 All living systems e.g. human being, plants, animals.



Closed system : This type of system can exchange energy, (in the form of heat, work or radiations) but not matter with its surroundings. The boundary is sealed but not insulated. Amount of the system will remain constant.

Eg.1 Coffee in closed vessel, Hot water in closed vessel.

Eg.2 Glowing bulb, tube light.




Isolated system : This type of system are perfectly insulated systems and cannot interact in any way with its surrounding i.e. neither matter nor energy can be exchanged with the surrounding. The boundary is sealed and insulated. Universe can be considered as an isolated system.

Eg.1 Coffee in thermos flask.



State of the System :

- Properties which define state of any system are called its state variables or thermodynamic variables or thermodynamic quantities.



- The state of the system is defined by its measurable properties like temperature, pressure, volume etc.
- If any of these properties change, state of the system is said to be changed.

State Function :

Those state variables which depend only upon state of the system but doesn't depend upon the path or mechanism followed by the system to achieve final state are called state function.

State functions are denoted by capital letters.


Ex. E, H, S, G, T, P, V etc.

Path function :

Properties of the system which depend upon the initial and final state of the system as well as the path or mechanism followed by the system to achieve final state are called path function.

Path functions are denoted by small letters.

Eg. Work done (w), heat (q)



- (i) **Intensive properties** : The properties of the system which are independent of matter (size and mass) present in system are called intensive properties.
- (ii) **Extensive properties** : The properties of the system which are dependent on matter (size and mass) present in system are called extensive properties

Extensive Properties	Intensive Properties
Volume (V)	Molar volume (V_m)
Number of moles (n)	Density (d)
Mass (M)	Gibb's energy per mole (G_m)
Gibb's Energy (G)	Specific heat
Entropy (S)	Pressure (P)
Enthalpy (H)	Temperature (T)
Internal energy (E or U)	All concentration terms (M , N)
Heat capacity (C)	Boiling point, freezing point (T_b , T_f)
Force (F)	Cell potential (E_{cell})
Surface Area (A)	Specific conductance (k)
	Refractive index
	Surface tension, Viscosity
	pH value
	Vapour pressure

Special Points

(i) The ratio of two extensive properties indicates the intensive property

Eg. $d = \frac{\text{mass}}{\text{volume}}$

(ii) An extensive property can be converted into intensive property when it is defined for unit amount of the substance.

Eg. Mass per unit volume = density; $\left(d = \frac{m}{V}\right)$

(iii) Intensive properties of a substance are non-additive in nature while extensive properties are additive in nature.

Vessel	H ₂ O A	H ₂ O B	On adding
Mass	m ₁	m ₂	m ₁ + m ₂ (Total mass)
Moles	n ₁	n ₂	n ₁ + n ₂ (Total moles)
Volume	V ₁	V ₂	V ₁ + V ₂ (Total volume)
Density	d	d	Remains same (d)
Boiling point	T	T	Remains same (T)


Extensive

Intensive

Types of Thermodynamic Processes :

When a system changes from one state to another, the operation is called a thermodynamic process.

Thermodynamic processes may be in form of expansion or compression.



(a) Isothermal Process (n, T = constant)

Isothermal processes are those processes in which

(a) Temperature of system during entire process remains constant i.e. (dT = 0)

(b) Heat is exchanged with surroundings

(c) Volume and pressure are variable

For ideal gas in Isothermal process $\Delta E = 0$ and $\Delta H = 0$

All phase transitions are isothermal process but $\Delta E \neq 0$ and $\Delta H \neq 0$

(b) Isobaric Process ($n, P = \text{constant}$) :

Isobaric processes are those processes in which

- (a) Pressure of system during entire process remains constant i.e. $dP = 0$
- (b) Volume and temperature are variable.
- (c) Process in open system is isobaric in nature.

(c) Isochoric Process ($n, V = \text{constant}$) :

Isochoric processes are those processes in which

- (a) Volume remains constant i.e. $dV = 0$
- (b) Pressure and temperature are variable
- (c) Work, $w = -P\Delta V$, $\therefore \Delta V = 0$, $\therefore w = 0$ (Zero)
- (d) Process in closed system is isochoric in nature.

(d) Adiabatic Process ($n = \text{constant}, q = 0$)


Adiabatic processes are those processes in which

- (a) No exchange of heat between system and surrounding takes place during entire process i.e. $q = 0$
- (b) The temperature, pressure, volume of the system varies.
- (c) The system is thermally insulated by keeping the system in an insulated container.

(e) Cyclic Process :

When a system undergoes a number of different processes and finally returns to its initial state, it is termed as cyclic process.

In cyclic process change in all state function will be zero. i.e. $\Delta E = 0, \Delta H = 0, \Delta P = 0, \Delta T = 0$



Reversible Process (quasi-static) :

- (1) Process in which all changes occurring at any part of the system are exactly reversed when small changes in variables are carried out in opposite direction.
- (2) Driving force should be infinitesimally greater than opposing force.
- (3) Process takes place in infinitesimal small steps or in many steps and takes infinite time to complete the process.
- (4) It is an ideal process.
- (5) Work obtained in expansion is maximum.
- (6) System is in virtual equilibrium at any state.
- (7) $P_{\text{ext}} = P_{\text{int}} \pm dP$; P_{ext} is variable.

Irreversible Process :

- (1) Process in which direction of change cannot be reversed by small changes in variables.
- (2) Driving force is much greater than opposing force.
- (3) It takes finite time and finite/usually single step.
- (4) Process takes place in short time
- (5) All natural processes are irreversible
- (6) System is in equilibrium only at initial and final state
- (7) $P_{\text{ext}} = P_{\text{int}} \pm \Delta P$; P_{ext} is constant.

Sign Conventions :

The work (w) is positive, when work is done on the system and w is negative, when the work is done by the system. The heat (q) is positive, when heat is transferred from surroundings to the system and q is negative when heat is transferred from system to the surroundings.

Sign convention

If 'q' amount of heat is given to the system


$$q = +ve$$



If 'w' amount of work is done on the system

$$w = +ve$$

(Compression)



If 'q' amount of heat is released /evolved by the system

$$q = -ve$$



If 'w' amount of work is done by the system

$$w = -ve$$

(Expansion)

**Work and Heat :**

Work : Product of force and displacement is known as work.

$$\text{work (w)} = \text{force (F)} \times \text{displacement (}\ell\text{)}$$

Consider a gas enclosed in a cylinder fitted with a frictionless piston.

Suppose area of cross section of cylinder = A and pressure on the piston = P

Initial volume of the gas = V_1 and final volume of the gas = V_2

(By expansion) displacement of piston = ℓ

$$\text{work done by the gas (in expansion)} = w = F \cdot \ell$$

$$\therefore P = \frac{F}{A} \quad \therefore F = P \times A$$

$$w = P \times A \times \ell \quad (\text{change in volume} = A \times \ell = V_2 - V_1)$$

$$w = P \times (V_2 - V_1)$$

$$w = P\Delta V \quad (\text{According to Physics})$$

In general

$$w = -P_{\text{external}} \Delta V \quad (\text{According to Chemistry})$$


Units of heat & work :

Calorie: It is defined as the quantity of heat required to raise the temperature of 1 g of water by 1°C (14.5 to 15.5°C)

$$1 \text{ Cal} = 4.184 \text{ J} \simeq 4.2 \text{ J}$$

$$1 \text{ L-atm} = 101.3 \text{ J} = 24.206 \text{ Cal} = 101.3 \times 10^7 \text{ erg}$$

$$1 \text{ L-atm} > 1 \text{ Cal} > 1 \text{ J} > 1 \text{ erg}$$



★ **Golden Key Points** ★

- Molar properties like H_m , G_m , S_m , U_m are intensive properties.
- In cyclic process change in all state function will be equal to zero.
 $\Delta E = 0$; $\Delta H = 0$, $\Delta P = 0$, $\Delta T = 0$ etc.
- For ideal gases $\Delta E = 0$ (For isothermal process)
- All natural process are irreversible in nature.

Illustration 1:

Find the work in each case :

(a) When one mol of ideal gas in 10 litre container at 1 atm is allowed to enter a vacuum bulb of capacity 100 litre.

(b) When 1 mol of gas expands from 1 litre to 5 litre against constant atmospheric pressure.

Solution:

(a) $W = -P\Delta V$

but since gas enters the vacuum bulb and pressure in vacuum is zero. This type of expansion is called **free expansion** and work is zero.

Note :- Work in free expansion is always zero.

(b) $W = -P\Delta V = -1(5 - 1) = -4 \text{ L-atm}$.

Illustration 2:

A 5 litre cylinder contained 10 mol of oxygen gas at 27°C . Due to sudden leakage through the hole, all the gas escaped into the atmosphere and the cylinder got empty. If the atmosphere pressure is 1.0 atm. Find work (in L atm) ($R = 0.083 \text{ L atm mol}^{-1} \text{ K}^{-1}$)

Solution:

$$V_{\text{initial}} = 5 \text{ L}$$

$$T = 27^\circ\text{C} = 27 + 273 = 300 \text{ K}$$

$$V_{\text{final}} = \frac{nRT}{P} = \frac{10 \times 0.083 \times 300}{1} = 249 \text{ L}$$

$$\Delta V = V_{\text{final}} - V_{\text{initial}}$$

$$= 249 - 5 = 244 \text{ L}$$

$$W_{\text{exp}} = -P\Delta V$$

$$= -1 \times 244 \text{ L-atm}$$

$$= -244 \text{ L-atm}$$



BEGINNER'S BOX-1

- Which of the following is open system
(1) Animals and plants (2) A fridge (3) A solar cooker (4) None of these
CTD226
- One mole of gas occupying 3 litre volume is expanded against a constant external pressure of one atm to a volume of 15 litre. The work in this process is :-
(1) -1.215×10^3 J (2) $+12.15 \times 10^3$ J (3) $+121.5 \times 10^3$ J (4) $+1.215 \times 10^3$ J
CTD227
- The work during the expansion of a gas from a volume of 4 dm³ to 6 dm³ against a constant external pressure of 3 atm is :-
(1) -608 J (2) +304 J (3) -304 J (4) -6 J
CTD228
- The work during the expansion of a gas from a volume of 14 dm³ to 16 dm³ against a constant external pressure of 2 atm is :-
(1) -405.2 J (2) +304 J (3) -304 J (4) -6 J
CTD229

Internal Energy (E/U) :

Internal energy of a system is defined as the sum of different energies associated with its atoms and molecules like Potential energy, Kinetic energy (due to translatory, rotatory and vibratory motion), electronic energy, nuclear energy etc.

$$E = E_{PE} + E_T + E_R + E_V + E_e + E_N$$

(i) Internal energy is an **extensive property**.

(ii) We can **never find out the absolute value** of internal energy (E) of system.

We can only calculate the change in internal energy of the system (ΔE) by using an instrument which is called as Bomb calorimeter. In Bomb calorimeter reactions are carried out at constant volume.

(iii) Internal energy is a **state function**.

$$\Delta E = E_f - E_i$$

(In reaction) $\Delta E = E_p - E_R$

ΔE is + ve if $E_f > E_i$

ΔE is - ve if $E_f < E_i$

Laws of Thermodynamics :**General Point :**

Laws of thermodynamics are based on human experiences and there is no formal proof for them.

First Law of Thermodynamics (FLOT) :

-  This law is based on law of conservation of energy and was given by **Robert Mayer and Helmholtz**.
-  Energy can neither be created nor destroyed but can be transformed from one form to another.
-  The total energy of the universe is always constant. i.e. total energy of an isolated system is always conserved.
-  The mathematical form of first law of thermodynamics can be expressed as follows

$\Delta E = q + w$: where q , w are path function and E is state function.

Here q is the energy given to the system and w is the work done on the system; ΔE is change in internal energy.

Illustration 3:

1g of water changes from liquid to vapour phase at constant pressure of 1 atmosphere, the volume increases from 1 mL to 1671 mL. The heat of vaporisation at this pressure is 540 Cal g⁻¹ Find the increase in internal energy of water. (1 L atm = 101 J)

Solution:

Work done

$$w = -P\Delta V = -P(V_2 - V_1)$$

$$= -1(1671 - 1) \times \frac{1}{1000} = \frac{-1670}{1000} \text{ L-atm}$$

$$= \frac{-1670}{1000} \times 101 \text{ J} = -168.67 \text{ J}$$

$$\text{given that } q = 540 \text{ Cal} = 540 \times 4.2 \text{ J} = 2268 \text{ J}$$

$$\therefore \Delta E = q + w = 2268 - 168.67 = 2099.33 \text{ J}$$

Illustration 4:

A gas occupies 2 L at STP. It is provided 300 J heat so that its volume becomes 2.5 L at 1 atm. Calculate change in its internal energy.

Solution:

$$w = -P\Delta V = -1 \times (2.5 - 2) = -0.5 \text{ L-atm or } w = -0.5 \times 101.3 = -50.65 \text{ J.}$$

$$\Delta E = q + w = 300 + (-50.65)$$

$$\Delta E = 249.35 \text{ J}$$

Illustration 5:

A sample of gas present in a cylinder fitted with a frictionless piston expands against a constant pressure of 1 atm from a volume of 2L to 12L. During the process, it absorbs 600 J of heat from the surroundings. Calculate the change in internal energy of the system.

Solution:

During the process,

$$q = 600 \text{ J}, \Delta V = 12 - 2 = 10 \text{ L}, P = 1 \text{ atm}$$

$$w = -P\Delta V$$

$$= -1 \times 10 = -10 \text{ L atm}$$

$$\text{Now, } 1 \text{ L atm} = 101.3 \text{ J}$$

$$\therefore w = -10 \times 101.3 = -1013 \text{ J}$$

According to first law of thermodynamics,

$$\Delta E = q + w = 600 - 1013 = -413 \text{ J}$$

Illustration 6:

Two moles of an ideal gas at 2 atm and 27°C is compressed isothermally to one half of its volume by a constant external pressure of 4 atm. Calculate q , w & ΔE . ($R = 0.082 \text{ L atm mol}^{-1} \text{ K}^{-1}$)

Solution:

$$w = -P_{\text{ext}}\Delta V = -P_{\text{ext}}(V_f - V_i)$$

$$V_i = \frac{nRT}{P}$$

$$n = 2 \text{ mol and } R = 0.082 \text{ atm L mol}^{-1} \text{ K}^{-1},$$

$$T = 273 + 27 = 300 \text{ K}, P = 2 \text{ atm}$$

$$V_i = \frac{2 \times 0.082 \times 300}{2} = 24.6 \text{ L and } V_f = \frac{V_i}{2} = \frac{24.6}{2} = 12.3 \text{ L}$$

$$w = -4 \text{ atm} \times (12.3 - 24.6) \text{ L}$$

$$= 49.2 \text{ L atm}$$

$$= 49.2 \times 101.3 \text{ J} = 4984 \text{ J.}$$

Since, it is isothermal compression, $\Delta E = 0$

$$\text{Now, } \Delta E = q + w$$

$$0 = q + 4984 \text{ J or } q = -4984 \text{ J}$$

Illustration 7:

A system is provided with 100 J of heat. Work done on the system is 20 J. What is the change in internal energy ?

Solution:

$$q = 100 \text{ J} \quad \text{and} \quad w = +20 \text{ J}$$

$$\Delta E = q + w$$

$$= 100 + 20 \text{ J} = 120 \text{ J}$$

Illustration 8:

An insulated container is divided into two equal portions. One portion contains an ideal gas at pressure P and temperature T , while the other portion is a perfect vacuum. If a hole is opened between the two portions, Calculate the –

(i) Change in internal energy of the gas

(ii) Change in temperature of the gas

Solution:

For insulated system, $q = 0$

The gas is allowed to expand against vacuum, the process of free expansion and thus

$$w = -P\Delta V = 0 \quad (\text{for vacuum} = P_{\text{ext}} = 0)$$

Thus from first law of thermodynamics, $\Delta E = q + w$ or $\Delta E = 0$

i.e. internal energy change is zero or internal energy of gas remains constant during free expansion.

Also $E \propto T$ and thus temperature of the gas will also remain constant.

Conclusions from the First Law of Thermodynamics :**(a) During isothermal process of an ideal gas :**

During an isothermal process the temperature of the system remains constant and hence

$$\Delta E = 0$$

Therefore

$$\text{FLOT : } \Delta E = q + w$$

$$\therefore +q = -w \quad \text{or} \quad -q = +w$$

In isothermal process –

(I) Heat absorbed by the system is equal to work done by the system.

OR

(II) Heat evolved by the system is equal to work done on the system.

(b) During isochoric process :

At constant volume $V_1 = V_2$ i.e. $\Delta V = 0$

Hence $W = -P\Delta V = 0$ No work done at constant volume therefore,

$$\text{FLOT: } \Delta E = q + w$$

$$\Rightarrow \Delta E = q_v$$

(i) In isochoric process

- At constant volume, heat absorbed by the system is equal to increase in internal energy of the system.

OR

- At constant volume, heat evolved by the system is equal to decrease in internal energy of the system.

(ii) Heat at constant volume (q_v) = ΔE

(iii) **In isochoric process heat is independent of path.**

(c) During adiabatic process :

During adiabatic process the system acts as an isolated system and hence $q = 0$ in such cases.

Therefore

$$\text{FLOT : } \Delta E = q + w \quad \because q = 0 \quad \Delta E = w$$

(i) Work done on the system is equal to increase in internal energy of the system i.e., when a gas is compressed adiabatically its internal energy increases.

OR

Work done by the system is equal to decrease in internal energy of the system, i.e., when a gas is expanded adiabatically its internal energy decreases.

(ii) **In adiabatic process work is independent of path.**

(d) During Cyclic Process :

$$\Delta E = 0 \quad \text{FLOT ; } \Delta E = q + w \quad \Rightarrow q = -w$$

In cyclic process.

(i) Work done by the system is equal to heat absorbed by the system.

OR

(ii) Work done on the system is equal to heat evolved by the system.

(e) During Isobaric process :

P is constant

$$\text{FLOT : } \Delta E = q + w$$

$$\Delta E = q + (-P\Delta V)$$

$$q_p = \Delta E + P\Delta V$$

$$q_p = (E_2 - E_1) + P(V_2 - V_1)$$

$$q_p = (E_2 + PV_2) - (E_1 + PV_1)$$

$$\because \text{Enthalpy } H = E + PV$$

$$\therefore q_p = H_2 - H_1$$

$$q_p = \Delta H$$

(i) **In Isobaric process :**

- At constant pressure, heat absorbed by the system is equal to increase in enthalpy of the system.

OR

- At constant pressure, heat evolved by the system is equal to decrease in enthalpy of the system.

(ii) Heat at constant pressure (q_p) = ΔH

(iii) **In isobaric process heat is independent of path.**



BEGINNER'S BOX-2

- A system absorb 300 cal of heat with the result of that, the volume of the system becomes double of its initial volume and temperature changes from 273K to 546K. The work done by the system on the surroundings is 200.0 Cal. Calculate ΔE .
 (1) 273 Cal (2) 500 Cal (3) 100 Cal (4) -500 Cal
CTD230
- One mol of an ideal gas at 300 K is expanded isothermally from an initial volume of 1 litre to 10 litre. The ΔE for the process is :- ($R = 2 \text{ Cal K}^{-1} \text{ mol}^{-1}$)
 (1) 163.7 Cal (2) 1381.1 Cal (3) 9 L-atm (4) Zero
CTD231
- In an adiabatic process, which of the following is true ?
 (1) $q = +w$ (2) $-\Delta E = -w$ (3) $P\Delta V = 0$ (4) $q = \Delta E$
CTD232
- In an isochoric process, the increase in internal energy is :-
 (1) Equal to the heat absorbed (2) Equal to the heat evolved
 (3) Equal to the work done (4) Equal to zero
CTD233

Enthalpy (H) :

Mathematically it is heat contained in the system measured at constant pressure.

The sum of internal energy and pressure volume (PV) energy is known as enthalpy.

$$\Delta H = \Delta E + \Delta(PV)$$



It is impossible to determine absolute value of enthalpy, so, we determine change in enthalpy (ΔH).

$$\Delta H = H_{\text{final}} - H_{\text{initial}}$$



Enthalpy is an extensive property because E and V are extensive properties.



It is a state function because E, P and V are state functions.

$$\therefore H = E + PV$$

$$\therefore \Delta H = \Delta E + \Delta(PV) \quad \dots(i)$$

(when P, V and T are variables)

$$\text{At constant pressure :} \quad \Delta H = \Delta E + P.\Delta V \quad \dots(ii)$$

$$\text{At constant volume :} \quad \Delta H = \Delta E + V.\Delta P \quad \dots(iii)$$

For chemical reactions at constant temperature and pressure

$$\therefore P.\Delta V = \Delta n_g RT$$


$$\text{So from equation (i)} \quad \Delta H = \Delta E + \Delta n_g RT \quad \dots(iv)$$

where $\Delta H = q_p$; at constant P ; $\Delta E = q_v$; at constant V

$$\text{So equation (iv) can be also written as} \quad q_p = q_v + \Delta n_g RT \quad \dots(v)$$

★ Golden Key Points ★

- If, $\Delta n_g = 0 \rightarrow \Delta H = \Delta E$ eg. $\text{H}_2(\text{g}) + \text{I}_2(\text{g}) \rightarrow 2\text{HI}(\text{g})$
- If, $\Delta n_g > 0 \rightarrow \Delta H > \Delta E$ eg. $\text{PCl}_5(\text{g}) \rightarrow \text{PCl}_3(\text{g}) + \text{Cl}_2(\text{g})$
- If, $\Delta n_g < 0 \rightarrow \Delta H < \Delta E$ eg. $\text{N}_2(\text{g}) + 3\text{H}_2(\text{g}) \rightarrow 2\text{NH}_3(\text{g})$

**Illustration 9:**

The heat of reaction for $\text{C}_{10}\text{H}_8(\text{s}) + 12\text{O}_2(\text{g}) \longrightarrow 10\text{CO}_2(\text{g}) + 4\text{H}_2\text{O}(\ell)$ at constant volume is -1228.2 kcal at 25°C . Calculate the heat of reaction at constant pressure and at 25°C .

Solution:

$$\Delta n_g = [10] - [12] = -2$$

$$\Delta H = \Delta E + \Delta n_g RT$$


$$= -1228.2 \times 10^3 + (-2)(2) \times 298$$

$$= -1229392 \text{ Cal}$$

$$\Delta H = -1229.392 \text{ kCal}$$

Illustration 10:

For the reaction at 25°C



Calculate ΔE° of the reaction at the given temperature.

Solution:

$$\Delta H^\circ = \Delta E^\circ + \Delta n_g RT$$

$$\Delta n_g = 2 - 1 = 1 \text{ mol}$$

$$\Delta E^\circ = \Delta H^\circ - \Delta n_g RT$$

$$= 11.04 \text{ Kcal} - 1 \text{ mol} \times \frac{2}{1000} \text{ kCal mol}^{-1} \text{K}^{-1} \times 298 \text{K}$$

$$= 11.04 - 0.596 = 10.44 \text{ kCal}$$

Illustration 11:

At 27°C the internal energy change of reaction $\text{H}_2(\text{g}) + \text{Cl}_2(\text{g}) \longrightarrow 2\text{HCl}(\text{g})$ is 2Cal. What is the enthalpy change of this reaction ?

Solution:

$$\Delta H = \Delta E + \Delta n_g RT$$

$$\Delta H = \Delta E + 0 \times RT$$

$$\Delta H = \Delta E$$


$$\Delta H = 2\text{Cal}$$

Illustration 12:

The heat of combustion of gaseous methane (CH_4) at constant volume is measured in bomb calorimeter at 298K is found to be $-885.4 \text{ kJ mol}^{-1}$. Find the value of enthalpy change at the same temperature.

Solution:

Combustion of methane gives $\text{CO}_2(\text{g})$ and $\text{H}_2\text{O}(\ell)$ as



$$\Delta E = -885.4 \text{ kJ mol}^{-1} = -885400 \text{ J mol}^{-1}$$

$$\Delta n_g = 1 - (1 + 2) = -2 \text{ mol}$$

$$T = 298\text{K}, R = 8.314 \text{ J mol}^{-1} \text{ K}^{-1}$$

$$\text{Now, } \Delta H = \Delta E + \Delta n_g RT$$

$$= -885400 + (-2 \text{ mol}) \times (8.314 \text{ J mol}^{-1} \text{ K}^{-1}) \times (298\text{K})$$

$$= -885400 - 4955$$

$$= -890355 = -890.355 \text{ kJ}$$

Illustration 13:

The enthalpy change (ΔH) for the reaction : $\text{N}_2(\text{g}) + 3\text{H}_2(\text{g}) \rightarrow 2\text{NH}_3(\text{g})$ is -92.38 kJ at 298 K. What is ΔE at 298 K ?

Solution:

$$\Delta H \text{ and } \Delta E \text{ are related as } \Delta H = \Delta E + \Delta n_g RT$$

$$\text{for the reaction, } \text{N}_2(\text{g}) + 3\text{H}_2(\text{g}) \rightarrow 2\text{NH}_3(\text{g})$$

$$\Delta n_g = 2 - (1 + 3) = -2 \text{ mol}, T = 298 \text{ K}$$

$$\Delta H = -92.38 \text{ kJ} = -92380 \text{ J}, R = 8.314 \text{ J K}^{-1} \text{ mol}^{-1}$$

$$-92380 = \Delta E + (-2 \text{ mol}) \times (8.314 \text{ J mol}^{-1} \text{ K}^{-1}) \times (298 \text{ K})$$

$$-92380 = \Delta E - 4955$$

$$\Delta E = -92380 + 4955$$

$$= -87425 \text{ J} = -87.425 \text{ kJ}.$$

Illustration 14:

The enthalpy change for the reaction $\text{CaCO}_3(\text{s}) \rightarrow \text{CaO}(\text{s}) + \text{CO}_2(\text{g})$ at 1000K is 176 kJ mol^{-1} . Calculate the change in internal energy.

Solution:

$$\Delta H = \Delta E + \Delta n_g RT$$

$$176 = \Delta E + (+1) \times 8.314 \times 10^{-3} \times 1000$$


$$\Delta E = 167.686 \text{ kJ}$$



BEGINNER'S BOX-3

- When a solid melts, there is :-
 (1) No increase in enthalpy (2) Increase in enthalpy
 (3) Decrease in enthalpy (4) Anything can happen
 CTD234
- For the reaction $\text{N}_2(\text{g}) + 3\text{H}_2(\text{g}) \longrightarrow 2\text{NH}_3(\text{g})$ which of the following is valid ?
 (1) $\Delta H = \Delta E$ (2) $\Delta H < \Delta E$ (3) $\Delta H > \Delta E$ (4) None of these
 CTD235
- Heat exchanged in a chemical reaction at constant pressure is called as :-
 (1) Internal energy (2) Enthalpy (3) Entropy (4) Free energy
 CTD236
- Latent heat of vaporisation of a liquid at 500 K and 1 atm pressure is 10.0 kCal mol⁻¹. What will be the change in internal energy of 3 mol of liquid at same temperature and pressure ?
 (1) 13.0 kCal (2) -13.0 kCal (3) 27.0 kCal (4) -27.0 kCal
 CTD237
- What is the value of Δn_g if we consider the combustion of 1 mol of liquid ethanol if reactants and products are at 298 K ?
 (1) -1 (2) -2 (3) +1 (4) +2
 CTD238
- If a reaction involves only solids and liquids, which of the following is true ?
 (1) $\Delta H < \Delta E$ (2) $\Delta H = \Delta E$ (3) $\Delta H > \Delta E$ (4) $\Delta H = \Delta E + RT\Delta n_g$
 CTD239
- The value of $\Delta H - \Delta E$ for the following reaction at 27°C will be, $2\text{NH}_3(\text{g}) \longrightarrow \text{N}_2(\text{g}) + 3\text{H}_2(\text{g})$:-
 (1) $8.314 \times 273 \times (-2)$ J (2) $8.314 \times 300 \times (-2)$ J
 (3) $8.314 \times 27 (+2)$ J (4) $8.314 \times 300 (+2)$ J
 CTD240
- At constant temperature for the reaction $\text{C}_3\text{H}_8(\text{g}) + 5\text{O}_2(\text{g}) \longrightarrow 3\text{CO}_2(\text{g}) + 4\text{H}_2\text{O}(\ell)$, $\Delta E - \Delta H$ is :-
 (1) +RT (2) -3RT (3) +3RT (4) -RT
 CTD241

Heat Capacity / Molar Heat Capacity / Specific Heat Capacity :



- (i) **Heat capacity (C)** : Amount of heat required to raise the temperature of given amount of a substance by 1°C or 1K is called heat capacity.

$$\text{Heat capacity} = \frac{\text{Heat required}}{\text{rise in temp.}}$$

$$C = \frac{q}{\Delta T}$$

Units :- JK⁻¹, Cal K⁻¹, J °C⁻¹, Cal °C⁻¹

- Heat capacity is extensive property.

- (ii) **Molar heat capacity (C_m)** : Amount of heat required to raise the temperature of 1 mole of substance by 1°C or 1 K is called as molar heat capacity.

$$\text{Molar heat capacity} = \frac{\text{Heat capacity}}{\text{mole of substance}} \Rightarrow C_m = \frac{C}{n}$$

Units :- $\text{J mol}^{-1} \text{ K}^{-1}$, $\text{Cal mol}^{-1} \text{ K}^{-1}$, $\text{J mol}^{-1} ^\circ\text{C}^{-1}$, $\text{Cal mol}^{-1} ^\circ\text{C}^{-1}$

- Molar heat capacity is an intensive property.

- (iii) **Specific heat capacity (c)** : Amount of heat required to raise the temperature of 1 g of substance by 1°C or 1 K is called as specific heat capacity.

$$C = \frac{C_m}{\text{molecular weight}}$$

Units :- $\text{J g}^{-1} \text{ K}^{-1}$, $\text{Cal g}^{-1} \text{ K}^{-1}$, $\text{J g}^{-1} ^\circ\text{C}^{-1}$, $\text{Cal g}^{-1} ^\circ\text{C}^{-1}$

- Specific heat capacity is an intensive property.
- If heat is supplied at constant pressure, then $C_p = \left(\frac{dq}{dT} \right)_p = \frac{dH}{dT} \quad \dots(i)$
- If heat is supplied at constant volume, then $C_v = \left(\frac{dq}{dT} \right)_v = \frac{dE}{dT} \quad \dots(ii)$

From equation (i) and (ii) :

From equation (i)	Unit	From equation (ii)
$\Delta H = C_p dT$ [Here C_p is heat capacity at constant P]	J K^{-1}	$\Delta E = C_v dT$ [Here C_v is heat capacity at constant V]
For n moles $\Delta H = nC_{p,m} dT$ [Here $C_{p,m}$ is molar heat capacity at constant p]	$\text{J mol}^{-1} \text{ K}^{-1}$	$\Delta E = nC_{v,m} dT$ [Here $C_{v,m}$ is molar heat capacity at constant V]
For m gram $\Delta H = mC_p dT$ [Here C_p is gram specific heat (specific heat capacity) at constant P]	$\text{J g}^{-1} \text{ K}^{-1}$	$\Delta E = mC_v dT$ [Here C_v is gram specific heat (specific heat capacity) at constant V]

Relation between C_p and C_v for 1 mole of an ideal gas :

$$\therefore H = E + PV$$

for ideal gas,

$$PV = nRT$$

$$PV = RT \text{ for 1 mole}$$

$$\therefore H = E + RT \text{ differentiate w.r.t. temperature}$$

$$\left(\frac{dH}{dT} \right) = \left(\frac{dE}{dT} \right) + R$$

$$C_{p,m} = C_{v,m} + R$$

(i) $C_{p,m} - C_{v,m} = R$ Mayer's formula

(ii) $\frac{C_p}{C_v} = \gamma$, γ = Poisson's ratio

Work Done in Different Process :**(a) Work done in reversible isothermal process :**

$$P_{\text{external}} = \text{Variable}$$

For reversible process

$$P_{\text{ext}} = P_{\text{gas}} \pm dP$$

$$\therefore w_{\text{rev.}} = -\int_{V_1}^{V_2} P_{\text{ext.}} dV$$

$$w_{\text{rev.}} = -\int_{V_1}^{V_2} (P_{\text{gas}} \pm dP) \cdot dV$$

Both dP and dV are very small so, $(dP \cdot dV)$ is negligible.

$$w_{\text{rev.}} = -\int_{V_1}^{V_2} P_{\text{gas}} \cdot dV$$

$$w_{\text{rev.}} = -\int_{V_1}^{V_2} \frac{nRT}{V} \cdot dV$$

$$w_{\text{rev.}} = -nRT \int_{V_1}^{V_2} \frac{dV}{V}$$

$$w_{\text{rev.}} = -nRT \ln \left(\frac{V_2}{V_1} \right)$$

or, $w_{\text{rev.}} = -2.303 nRT \log_{10} \left(\frac{V_2}{V_1} \right) \quad \dots(i)$

according to Boyle's law at constant temperature $P \propto \frac{1}{V}$ or $PV = \text{constant}$

$$\therefore P_1 V_1 = P_2 V_2 \quad \Rightarrow \quad \frac{V_2}{V_1} = \frac{P_1}{P_2}$$

$$w_{\text{rev.}} = -2.303 nRT \log_{10} \left(\frac{P_1}{P_2} \right) \quad \dots(ii)$$

(b) Work done in reversible adiabatic process :

In adiabatic process $q = 0$

$$\text{FLOT : } \Delta E = q + w$$

$$\Rightarrow w = \Delta E \Rightarrow w = nC_{V,m} \Delta T$$

$$w = nC_{V,m} (T_2 - T_1) \quad \dots(i)$$

$$\therefore C_{P,m} - C_{V,m} = R$$

$$\frac{C_P}{C_V} - 1 = \frac{R}{C_V}$$

$$\gamma - 1 = \frac{R}{C_V} \left(\because \gamma = \frac{C_P}{C_V} \right) \Rightarrow C_V = \frac{R}{\gamma - 1} \quad \dots(ii)$$

From equation (i) and (ii) $w = \frac{nR}{\gamma - 1} (T_2 - T_1) = w = \frac{P_2 V_2 - P_1 V_1}{(\gamma - 1)}$

State equations of reversible adiabatic process are :

$$PV^\gamma = \text{constant}$$

$$TV^{\gamma-1} = \text{constant}$$

$$T^\gamma P^{1-\gamma} = \text{constant}$$

★ Golden Key Points ★

- Work in rigid vessel is zero because volume remains constant.
- Work during free expansion of an ideal gas in vacuum is zero, because P_{ext} is zero.
- Work in a chemical reaction $w = -P\Delta V = -\Delta n_g RT$

Illustration 15:

5 moles of oxygen are heated at constant volume from 10°C to 20°C. What will be the change in the internal energy of gas? The molar heat capacity of oxygen at constant pressure,

$$C_p = 7.03 \frac{\text{Cal}}{\text{mol K}} \text{ and } R = 2 \text{ Cal mol}^{-1} \text{ K}^{-1}$$

Solution:

We know mayer's relation is $C_{p,m} - C_{v,m} = R$

$$C_{v,m} = C_{p,m} - R = 7.03 - 2 = 5.03 \text{ Cal mol}^{-1} \text{ K}^{-1}$$

$$\text{As we know } \Delta E = nC_{v,m}\Delta T = 5 \times 5.03 \times 10 = 251.5 \text{ Cal}$$

Illustration 16:

At 27°C, one mole of an ideal gas compressed isothermally and reversibly from a pressure of 2 atm to 10 atm. Calculate ΔE and q in calorie.

Solution:

$$\text{For isothermal process } \Delta E = 0 \text{ and } w = -2.303 nRT \log_{10} \left(\frac{P_1}{P_2} \right)$$

$$w = -2.303 \times 1 \times 2 \times 300 \times \log \left(\frac{2}{10} \right)$$

$$w = +2.303 \times 600 \times \log 5$$

$$w = +2.303 \times 600 \times 0.699$$

$$w = +965.87 \text{ Cal}$$

For isothermal process

$$\therefore w = -q$$

$$\therefore q = -965.87 \text{ Cal}$$

Illustration 17:

A gas expands from 3 dm³ to 5 dm³ against a constant pressure of 3 atm. The work done during expansion is used to heat 10 mol of water of temperature 290 K. Calculate final temperature of water (if specific heat of water is 4.184 Jg⁻¹K⁻¹)

Solution:

Since work is done against constant P and thus, irreversible

$$\Delta V = 5 - 3 = 2 \text{ dm}^3 = 2 \text{ L}, P = 3 \text{ atm}$$

$$w = -P\Delta V = -3 \times 2 \text{ L atm} = -6 \times 101.3 \text{ J} = -607.8 \text{ Joule}$$

Now this work is used up in heating water

$$w = n \times C \times \Delta T$$

$$607.8 = 10 \times (4.184 \times 18) \times \Delta T$$

$$\Delta T = 0.81 = T_2 - T_1$$

$$\therefore \text{Final temperature} = T_1 + \Delta T = 290 + 0.81 = 290.81 \text{ K}$$

Illustration 18:

A sample of 3 mol of an ideal gas at 200K and 2 atm is compressed reversibly and adiabatically until the temperature reaches 250K, given that molar heat capacity is $27.5 \text{ J K}^{-1} \text{ mol}^{-1}$ at constant volume, calculate w .

Solution:

$$C_{v,m} = 27.5 \text{ J K}^{-1} \text{ mol}^{-1}$$

During reversible adiabatic process

$$w = nC_{v,m} (T_2 - T_1) = 3 \times 27.5 \times 50 = 4125 \text{ Joule}$$

Illustration 19:

10 moles of an ideal gas at 27°C and 10 atm. pressure occupying a volume of 24.6 L undergoes the following changes.

- (i) Isothermal & reversible expansion to 246 L
- (ii) Isothermal and irreversible expansion to 246 L.
- (iii) Isochoric heating to 177°C .

Calculate the work in each transformation in kJ.

Solution:

- (i) Work in isothermal reversible process

$$\begin{aligned} w &= -2.303 \times nRT \log \left(\frac{V_2}{V_1} \right) \\ &= -2.303 \times 10 \times 8.31 \times 300 \times \log \left(\frac{246}{24.6} \right) \\ &= -57413.79 \text{ J} = -57.41 \text{ kJ} \end{aligned}$$

- (ii) Work in isothermal irreversible expansion process

$$\begin{aligned} w &= -P (V_2 - V_1) = -10 (246 - 24.6) = -2214 \text{ L-atm} \\ &= -2214 \times 101.3 \text{ J} = -224.3 \text{ kJ} \end{aligned}$$

- (iii) Work in isochoric change

$$\text{Since, } \Delta V = 0 \quad \therefore w = 0$$

Illustration 20:

Find the work, when 2 mol of a gas expands isothermally from 5 dm^3 to 40 dm^3 against a constant external pressure of 2 atm at 298K. Also calculate w_{rev} for the change.

Solution:

- (i) $w = -P\Delta V$

$$w = -2 \times (40 - 5)$$

$$w = -70 \text{ L atm} = -70 \times 101.3 \text{ J}$$

$$w = -7091 \text{ J}$$

- (ii) $w = -2.303 nRT \log \left(\frac{V_2}{V_1} \right)$

$$w = -2.303 \times 2 \times 8.314 \times 298 \log \left(\frac{40}{5} \right)$$

$$w = -10.3 \times 10^3 \text{ J}$$



BEGINNER'S BOX-4

- Calculate w for the isothermal reversible expansion of 1 mol of an ideal gas from an initial pressure of 1.0 bar to a final pressure of 0.1 bar at a constant temperature of 273 K :-
 (1) -5227.2 J (2) $+5227.2 \text{ J}$ (3) -2257 J (4) $+2257 \text{ J}$
- When 229 J of energy is supplied as heat at constant pressure to 3 mol Ar(g), the temperature of the sample is increased by 2.55K. Calculate the molar heat capacity at constant volume :-
 (1) $30 \text{ kJ K}^{-1} \text{ mol}^{-1}$ (2) $30 \text{ J K}^{-1} \text{ mol}^{-1}$ (3) $21.7 \text{ J K}^{-1} \text{ mol}^{-1}$ (4) $21.7 \text{ kJ K}^{-1} \text{ mol}^{-1}$

CTD242

CTD243

Spontaneous Process and Non-Spontaneous Process

(i)
Spontaneous process

(ii)
Non-spontaneous process

(i) Spontaneous process :

- The process which has a natural tendency to occur in a particular direction either of its own or after proper initiation under the given set of conditions.
- All natural processes are ir-reversible and spontaneous processes. The natural processes take place on their own in one direction only.

(ii) Non-spontaneous process :

- The process which does not occur on its own in a particular direction i.e. a process which does not have a natural tendency to occur in a particular direction either of its own or after initiation is called as non-spontaneous process.
- Non-spontaneous process may be made to occur when energy from some external source is supplied continuously throughout the process.

Examples of spontaneous process that need no initiation :

- Ex.1** Flow of water from high level to low level.
 Flow of heat from hot body to cold body.
 Flow of charge from high potential to low potential.
 Flow of gas from high pressure to low pressure.

- Ex.2** Melting of ice at 25°C
 Evaporation of water at 25°C
 Dissolution of common salt in water.

- Ex.3** Mixing of different non reacting gases

Examples of spontaneous process that need initiation :

- Ex.1** Burning of fuel (coal, petrol)
 $\text{C(s)} + \text{O}_2 \rightarrow \text{CO}_2$
 $\text{CH}_4 + 2\text{O}_2 \rightarrow \text{CO}_2 + 2\text{H}_2\text{O}$

- Ex.2** Lightning of candle

Criteria for a process to be spontaneous :

- Tendency to attain minimum energy or maximum stability (Energy Factor).
- Tendency to attain maximum randomness (Entropy Factor)

Entropy (s) :

- (i) The thermodynamic quantity, which is used to measure **degree of randomness** or **disorderness** of the system is called as entropy.

Entropy (s) $\propto$ Randomness or disorderness

- (ii) More is the disorderness, higher is the entropy.
 (iii) The ratio of heat absorbed by the system in isothermal and reversible manner to the temperature at which heat is absorbed is equals to the change in entropy.

$$\Delta S = \frac{q_{\text{rev}}}{T}$$

Where q_{rev} = heat absorbed by the system in a reversible manner at the temperature T

Unit : J K⁻¹ or Cal K⁻¹

- (iv) $\Delta S = S_{\text{final}} - S_{\text{initial}}$
 If $S_{\text{final}} > S_{\text{initial}}$: Then ΔS = positive
 If $S_{\text{final}} < S_{\text{initial}}$: Then ΔS = negative
 (v) Entropy is an extensive property and state function.

Factors affecting entropy of system :

- (i) If $\Delta n_g > 0$ then $\Delta S > 0$
 If $\Delta n_g < 0$ then $\Delta S < 0$
 (ii) Physical state : $S_{\text{solid}} < S_{\text{liquid}} < S_{\text{gas}}$
 (iii) On increasing gaseous moles entropy increases.
 (iv) On increasing temperature, S will increase.

Ex. $\text{Fe(s)} \rightarrow \text{Fe(s)} : \Delta S = \text{positive}$
 300 K 400 K

- (v) On decreasing pressure, S increases.

Ex. $\text{N}_2(\text{g}) \rightarrow \text{N}_2(\text{g}) : \Delta S = \text{positive}$
 5 atm 2 atm

- (vi) Mixture :

$$\left. \begin{array}{l} \text{Solid + solid} \\ \text{liquid + liquid} \\ \text{gas + gas} \end{array} \right\} S \uparrow$$
Some famous or extra ordinary examples of entropy change :

- (i) Entropy of graphite > Entropy of diamond.
 (ii) $\text{NH}_4\text{Cl(s)} + \text{aq} \longrightarrow \text{NH}_4^+(\text{aq}) + \text{Cl}^-(\text{aq})$
 In this process NH_4^+ and Cl^- ions are free to move in solution where as they are not free to move in solid NH_4Cl .
 Hence ΔS is positive for this type of dissolution process.
 (iii) On addition of HCl in the aqueous solution of Ag^+ ions entropy decreases due to precipitation of AgCl .
 (iv) **On boiling of egg :** Denaturation of proteins occur. Thus entropy increases.

- (v) **Stretching of rubber** : During stretching of rubber band its long flexible macromolecules get uncoiled. The uncoiled form has more specific geometry and more ordered arrangement. Thus entropy decreases.

Entropy change in a chemical reaction $\Delta S = \sum S_{\text{Product}} - \sum S_{\text{Reactant}}$

Entropy change for a process : $\Delta S = nC_V \ln \left(\frac{T_2}{T_1} \right) + nR \ln \left(\frac{V_2}{V_1} \right)$ or $\Delta S = nC_P \ln \left(\frac{T_2}{T_1} \right) + nR \ln \left(\frac{P_1}{P_2} \right)$

Case I : For an ideal gas reversible isothermal process; $T_2 = T_1$

$$\Delta S = nR \ln \left(\frac{V_2}{V_1} \right) = nR \ln \left(\frac{P_1}{P_2} \right)$$

$$\Delta S = 2.303 nR \log \left(\frac{V_2}{V_1} \right) = 2.303 nR \log \left(\frac{P_1}{P_2} \right)$$

Case II : For an isochoric process $V_2 = V_1$

$$\Delta S = nC_V \ln \left(\frac{T_2}{T_1} \right) ; \Delta S = 2.303 nC_V \log \left(\frac{T_2}{T_1} \right)$$

For reversible adiabatic process : $q_{\text{rev}} = 0$, $\Delta S = \frac{q_{\text{rev}}}{T} = 0$, $S = \text{constant}$

Entropy remains constant so process is also known as **isoentropic process**.

Entropy change during phase transition :

(I) Entropy of fusion $[(\Delta S)_f]$:

The entropy change, when 1 mol solid changes into liquid at its melting point temperature.
solid $\rightleftharpoons$ liquid

$$(\Delta S)_f = \frac{\Delta H_{\text{fusion}}}{T}$$

(II) Entropy of vapourisation $[(\Delta S)_{\text{vap}}]$:

The entropy change, when 1 mol liquid changes into vapour at its boiling point temperature.
liquid $\rightleftharpoons$ vapour

$$(\Delta S)_{\text{vap}} = \frac{\Delta H_{\text{vap}}}{T}$$

(III) Entropy of sublimation $[(\Delta S)_{\text{sub}}]$:

The entropy change, when 1 mol solid changes into vapour at a particular temperature.
solid $\rightleftharpoons$ vapour

$$(\Delta S)_{\text{sub}} = \frac{\Delta H_{\text{sub}}}{T}$$

Total entropy change in reversible process :

In reversible process, at every step system and surroundings remain in thermal equilibrium with each other. Let a system, releases q heat to the surroundings at temperature T .

$$\Delta S_{\text{system}} = \frac{-q_{\text{system}}}{T} ; \quad \Delta S_{\text{surroundings}} = \frac{+q_{\text{system}}}{T}$$

$$\therefore \Delta S_{\text{system}} = \Delta S_{\text{system}} + \Delta S_{\text{surroundings}}$$

$$\Delta S_{\text{total}} = \frac{-q_{\text{system}}}{T} + \frac{q_{\text{system}}}{T} \Rightarrow \Delta S_{\text{total}} = 0$$

Total entropy change in irreversible process :

Let a system is at high temperature T_1 and surroundings are at low temperature T_2 .

Let q amount of heat is released by the system.

$$\Delta S_{\text{system}} = \frac{-q}{T_1}, \quad \Delta S_{\text{surroundings}} = \frac{+q}{T_2}$$

$$\therefore \Delta S_{\text{total}} = \Delta S_{\text{system}} + \Delta S_{\text{surroundings}} = \frac{-q}{T_1} + \frac{q}{T_2}$$

$$\therefore \Delta S_{\text{total}} = +ve \quad (\because T_1 > T_2)$$

Total entropy change for irreversible spontaneous process is always greater than zero.

Spontaneity of a process in terms of total entropy change :

If $\Delta S_{\text{total}} = +ve \Rightarrow$ Spontaneous process
 If $\Delta S_{\text{total}} = -ve \Rightarrow$ Non spontaneous process
 If $\Delta S_{\text{total}} = 0 \Rightarrow$ Process is at equilibrium

Illustration 21:

The enthalpy change for transition of liquid water to steam is 40.8 kJ mol^{-1} at 373K . Calculate ΔS for the process.

Solution:

The entropy change for the vapourization of water is given by $\Delta S = \frac{\Delta H_{\text{vap}}}{T}$

Given $\Delta H_{\text{vap}} = 40.8 \text{ kJ mol}^{-1} = 40.8 \times 1000 \text{ J mol}^{-1}$ and $T = 373 \text{ K}$

$$\Delta S = \frac{40.8 \times 1000 \text{ J mol}^{-1}}{373 \text{ K}} = 109.38 \text{ JK}^{-1} \text{ mol}^{-1}$$

Illustration 22:

Calculate the change in entropy for the fusion of 1 mol of ice. The melting point of ice is 273K and molar enthalpy of fusion of ice = 6 kJ mol^{-1}

Solution:

$$\Delta S_f = \frac{\Delta H_f}{T} = \frac{6 \times 10^3}{273} = 21.97 \text{ JK}^{-1} \text{ mol}^{-1}$$

Illustration 23:

The enthalpy of vapourisation of liquid diethyl ether $(C_2H_5)_2O$, is 26.0 kJ mol^{-1} at its boiling point (35.0°C). Calculate ΔS for conversion of :-

- (i) Liquid to vapour and
(ii) Vapour to liquid at 35°C

Solution:

$$(i) \Delta S_{\text{vap.}} = \frac{\Delta H_{\text{vap.}}}{T} = \frac{26 \times 10^3}{308} = 84.41 \text{ JK}^{-1}\text{mol}^{-1}$$

$$(ii) \Delta S_{\text{cond.}} = \frac{\Delta H_{\text{cond.}}}{T} = \frac{-26 \times 10^3}{308} = -84.41 \text{ JK}^{-1}\text{mol}^{-1}$$

Illustration 24:

Which of the following processes are accompanied by increase of entropy :

- (i) Dissolution of iodine in a solvent $\{I_2(s) \longrightarrow I_2(aq.)\}$
(ii) HCl is added to $AgNO_3$ and a precipitate of $AgCl$ is obtained.
(iii) A partition is removed to allow two gases to mix.

Solution:

Increase of entropy : (i) and (iii)

**BEGINNER'S BOX-5**

- In any natural process, occurring in the universe :-
(1) Entropy is conserved (2) Entropy increases
(3) Entropy decreases (4) Entropy remains unchanged
CTD244
- The most random state of H_2O system is :-
(1) Ice (2) $H_2O(l)$ at 80°C ; 1 atm
(3) Steam (4) $H_2O(l)$ at 25°C ; 1 atm
CTD245
- Change in entropy is negative for :-
(1) $Br_2(l) \longrightarrow Br_2(g)$ (2) $C(s) + H_2O(g) \longrightarrow CO(g) + H_2(g)$
(3) $M_2(g, 10 \text{ atm}) \longrightarrow M_2(g, 1 \text{ atm})$ (4) $Fe \text{ (at } 400\text{K)} \longrightarrow Fe \text{ (at } 300\text{K)}$
CTD246
- Total entropy change in spontaneous adiabatic process is :-
(1) Zero (2) < 0 (3) > 0 (4) None of these
CTD247
- 5 mole of an ideal gas expand reversibly from a volume of 8 dm^3 to 80 dm^3 at a temperature of 27°C . The change in entropy is :-
(1) 41.57 JK^{-1} (2) -95.73 JK^{-1} (3) 95.73 JK^{-1} (4) -41.57 JK^{-1}
CTD248
- The latent heat of vapourisation of water at 100°C is 540 Cal g^{-1} . Calculate the entropy increase when one mole of water at 100°C is evaporated :-
(1) $26 \text{ Cal K}^{-1} \text{ mol}^{-1}$ (2) $1.45 \text{ Cal K}^{-1} \text{ mol}^{-1}$ (3) $367 \text{ Cal K}^{-1} \text{ mol}^{-1}$ (4) $1.82 \text{ Cal K}^{-1} \text{ mol}^{-1}$
CTD249
- Calculate enthalpy of vapourisation per mole of ethanol. Given $\Delta S = 109.8 \text{ J K}^{-1} \text{ mol}^{-1}$ and B.P. of ethanol is 78.5°C :-
(1) Zero (2) $38.594 \text{ kJ mol}^{-1}$ (3) 3.85 kJ mol^{-1} (4) None of these
CTD250

Second Law of Thermodynamics (SLOT) :

- (i) It states about the direction of flow of heat
- (ii) All natural processes in universe are ir-reversible process or natural processes are spontaneous process.
- (iii) Due to spontaneous process entropy of universe is increasing continuously i.e. entropy of an isolated system increases.


$$(\Delta S)_T = +ve$$

or $(\Delta S)_T > 0$

or $(\Delta S)_{\text{system}} + (\Delta S)_{\text{surr.}} > 0$

Gibb's Energy (G or F) :

- Gibb's energy is defined at constant temperature and pressure to predict spontaneity of a process.
- Gibb's energy is a thermodynamic quantity which is used to measure the capacity of system to do useful work or Gibb's energy is that part of the total energy of system which can be converted into useful work.



Here $P_{2(g)}$ = External pressure acting from outside the piston.

$P_{(g)}$ = Pressure of gas inside the piston.

- The term Gibbs energy was introduced to explain criteria of spontaneity in terms of system.
- Since ; energy = useful work + randomness energy

$$H = G + TS$$

$$G = H - TS \quad \dots(i)$$

So the function that takes both enthalpy and entropy of system into account is called Gibbs energy.

- Gibbs energy is an extensive property and state function.
- Absolute value of G can't be measured but change can be measured, So we discuss (ΔG)

Gibbs Energy :

$$\Delta G = G_{\text{final}} - G_{\text{initial}}$$

From eq. (i) $\Delta G = \Delta H - T\Delta S$

Where ΔG = Change in Gibb's energy

ΔH = Change in enthalpy

ΔS = Change in entropy

Relation between Gibb's energy change and non expansion work or useful work :

$$\text{FLOT : } \Delta E = q + W$$

If, work is done by the system, then, $\Delta E = q - W$ (i)

According to Gibb's, system does both expansion and non expansion work.

$$\therefore W = W_{\text{expansion}} + W_{\text{non expansion}}$$

$$W = P\Delta V + W_{\text{non expansion}}$$

Put W in equation (i)

$$\Rightarrow \Delta E = q - (P\Delta V + W_{\text{non expansion}})$$

$$q = \Delta E + P\Delta V + W_{\text{non expansion}}$$

$$q = \Delta H + W_{\text{non expansion}} \quad (\because \Delta H = \Delta E + P\Delta V)$$

$$T\Delta S = \Delta H + W_{\text{non expansion}} \quad (\because \Delta S = q/T)$$

$$W_{\text{non expansion}} = T\Delta S - \Delta H \quad (\because \Delta G = \Delta H - T\Delta S)$$

$$\Rightarrow \boxed{W_{\text{non expansion}} = -\Delta G} \quad \text{or} \quad \boxed{W_{\text{useful}} = -\Delta G}$$

The decrease in Gibb's energy of system is equal to the non expansion work or useful work.

Relation in between ΔG of system and ΔS_{total} or Gibb's energy change and spontaneity :

$$\because \Delta S_{\text{total}} = \Delta S_{\text{system}} + \Delta S_{\text{surroundings}}$$

Let system releases heat at constant temperature T and pressure P.

$$\therefore q = q_p = \Delta H$$

$$\begin{aligned} & -q_{\text{system}} = +q_{\text{surroundings}} \\ \Rightarrow & -\Delta H_{\text{system}} = +\Delta H_{\text{surroundings}} \end{aligned} \quad \left. \vphantom{\begin{aligned} & -q_{\text{system}} = +q_{\text{surroundings}} \\ \Rightarrow & -\Delta H_{\text{system}} = +\Delta H_{\text{surroundings}} \end{aligned}} \right\}$$

$$\Delta S_{\text{surr.}} = \frac{\Delta H_{\text{surr.}}}{T}$$

$$\Delta S_{\text{surr.}} = \frac{-\Delta H_{\text{sys}}}{T}$$

$$\therefore \Delta S_{\text{total}} = \Delta S_{\text{sys.}} + \left(\frac{-\Delta H_{\text{sys}}}{T} \right)$$

$$T\Delta S_{\text{total}} = T\Delta S_{\text{sys.}} - \Delta H_{\text{sys.}}$$

$$T\Delta S_{\text{total}} = -(\Delta H_{\text{sys.}} - T\Delta S_{\text{sys.}})$$

$$T\Delta S_{\text{total}} = -\Delta G_{\text{sys.}}$$

$$\text{or} \quad \boxed{\Delta G_{\text{sys.}} = -T\Delta S_{\text{total}}}$$

$$(i) \quad \text{If, } \Delta S_{\text{total}} = +ve \Rightarrow \Delta G_{\text{system}} = -ve \Rightarrow \text{spontaneous process}$$

$$(ii) \quad \text{If, } \Delta S_{\text{total}} = -ve \Rightarrow \Delta G_{\text{system}} = +ve \Rightarrow \text{non spontaneous process}$$

$$(iii) \quad \text{If, } \Delta S_{\text{total}} = 0 \Rightarrow \Delta G_{\text{system}} = 0 \Rightarrow \text{process is at equilibrium.}$$

$\Delta_r H$	$\Delta_r S$	$\Delta_r G$	Description
–	+	–	Reaction spontaneous at all temperature
–	–	–	(at low T) Reaction spontaneous at low temperature
–	–	+	(at high T) Reaction nonspontaneous at high temperature
+	+	+	(at low T) Reaction nonspontaneous at low temperature
+	+	–	(at high T) Reaction spontaneous at high temperature
+	–	+	(at all T) Reaction nonspontaneous at all temperatures

Relationship between standard Gibbs energy change (ΔG°) and Equilibrium constant (K_{eq}) :-

For a reaction $m_1A + m_2B \rightleftharpoons n_1C + n_2D$

$$\Delta G = \Delta G^\circ + RT \ln Q$$

At equilibrium, $\Delta G = 0$ and $Q = K_{eq}$

$$0 = \Delta G^\circ + RT \ln K_{eq}$$

$$\therefore \Delta G^\circ = -RT \ln K_{eq} \quad \dots(i)$$

$$\text{or } \Delta G^\circ = -2.303 RT \log_{10} K_{eq} \quad \dots(ii)$$

from equation (i)

$$\ln K_{eq} = -\frac{\Delta G^\circ}{RT}$$

$$\therefore K_{eq} = e^{-\Delta G^\circ/RT} \quad \dots(iii)$$

Illustration 25:

For a certain reaction the change in enthalpy and change in entropy are $40.63 \text{ kJ mol}^{-1}$ and 100 JK^{-1} . What is the value of ΔG at 27°C and indicate whether the reaction is possible or not ?

Solution:

We know that :

$$\Delta G = \Delta H - T\Delta S$$

$$T = 27 + 273 = 300\text{K}$$

$$\Delta H = 40.63 \times 10^3 \text{ J mol}^{-1} = 40630 \text{ J mol}^{-1}$$

$$\Delta S = 100 \text{ JK}^{-1}$$

$$\Delta G = 40630 - 300 \times 100 = 40630 - 30000 = +10630 \text{ J}$$

Positive value of ΔG indicates that the reaction is not possible.

Illustration 26:

For a reaction at 25°C enthalpy change (ΔH) and entropy change (ΔS) are $-11.7 \times 10^3 \text{ J mol}^{-1}$ and $-105 \text{ J mol}^{-1} \text{ K}^{-1}$ respectively. Find out whether this reaction is spontaneous or not.

Solution:

$$\Delta G = \Delta H - T\Delta S$$


$$= -11700 - 298 \times (-105) = +19590 \text{ J}$$

$$\Delta G = +ve, \text{ so reaction is non-spontaneous.}$$

Illustration 27:

Calculate the equilibrium constant for the reaction given below at 400K.

If $\Delta H^\circ = 77.2 \text{ kJ mol}^{-1}$ and $\Delta S^\circ = 122 \text{ J K}^{-1} \text{ mol}^{-1}$

**Solution:**

$$\Delta G^\circ = \Delta H^\circ - T\Delta S^\circ = 77200 - 400 \times 122 = 28400 \text{ J}$$

$$\Delta G^\circ = -2.303 RT \log K_c$$

$$\text{or } 28400 = -2.303 \times 8.31 \times 400 \log K_c$$

$$\text{or } K_c = 1.958 \times 10^{-4}$$

Illustration 28:

For the reaction, $\text{N}_2(g) + 3\text{H}_2(g) \longrightarrow 2\text{NH}_3(g)$; $\Delta H = -95.4 \text{ kJ}$ and $\Delta S = -198.3 \text{ J K}^{-1}$. Calculate the temperature at which the reaction will proceed in forward direction.

Solution:

$$\Delta G = \Delta H - T\Delta S$$

$\therefore$ At equilibrium $\Delta G = 0$

$$\therefore \Delta H = T\Delta S$$

$$\text{so } T = \frac{-95.4 \times 1000 \text{ J}}{-198.3 \text{ J K}^{-1}} = 481 \text{ K}$$

For this reaction ΔH is -ve and ΔS is -ve, so it will be spontaneous at low temperature.

$\therefore$ Below 481K the reaction would be spontaneous.

Illustration 29:

Enthalpy and entropy changes of a reaction are $40.63 \text{ kJ mol}^{-1}$ and $108.8 \text{ J K}^{-1} \text{ mol}^{-1}$ respectively. Analyse the feasibility of the reaction at 27°C .

Solution:

$$\Delta H = 40.63 \text{ kJ mol}^{-1} = 40630 \text{ J K}^{-1} \text{ mol}^{-1}$$

$$\Delta S = 108.8 \text{ J K}^{-1} \text{ mol}^{-1}$$

$$T = 27^\circ\text{C} = 27 + 273 = 300 \text{ K}$$

$$\text{Now } \Delta G = \Delta H - T\Delta S$$

$$= 40630 \text{ J mol}^{-1} - (300 \text{ K}) \times (108.8 \text{ J K}^{-1} \text{ mol}^{-1})$$

$$\Delta G = 7990 \text{ J mol}^{-1}.$$

Since ΔG is positive, the reaction is not feasible in the forward direction.

Illustration 30:

For a certain reaction the change in enthalpy and change in entropy are $40.63 \text{ kJ mol}^{-1}$ and 100 J K^{-1} . Show that at 27°C the reaction is possible or not.

Solution:

$$\Delta G = \Delta H - T\Delta S$$

$$\Delta G = 40.63 - 300 \times 100 \times 10^{-3}$$

$$\Delta G = +10.630 \text{ kJ}$$

ΔG is positive so reaction is not possible.

Illustration 31:

Zinc reacts with dilute hydrochloric acid to give hydrogen at 17°C. The enthalpy of the reaction is $-12.55 \text{ kJ mol}^{-1}$ and entropy change is $5 \text{ JK}^{-1} \text{ mol}^{-1}$ for the reaction. Calculate the free energy change and predict whether the reaction is spontaneous or not.

Solution:

Given $\Delta H = -12.55 \text{ kJ mol}^{-1}$ and $\Delta S = 5 \text{ JK}^{-1} \text{ mol}^{-1}$

$T = 17 + 273 = 290 \text{ K}$

Applying $\Delta G = \Delta H - T\Delta S$

$$\begin{aligned} &= -12.55 \frac{\text{kJ}}{\text{mol}} - 290 \text{ K} \times \frac{5}{1000} \frac{\text{kJ}}{\text{K mol}} \\ &= -12.55 - 1.45 = -14 \frac{\text{kJ}}{\text{mol}} \end{aligned}$$

Since ΔG is negative, the reaction will be spontaneous

Illustration 32:

For a reaction both ΔH and ΔS are positive. Under what condition will the reaction occur spontaneously?

Solution:

The reaction will occur spontaneously only when $T\Delta S > \Delta H$.

$$\Delta G = \Delta H - T\Delta S = (+) - T(+)$$

For ΔG to be negative, $T\Delta S$ must be $> \Delta H$

Illustration 33:

Which of the following are state function ?

- (i) q (ii) Entropy (iii) Gibbs's energy (iv) H (v) w

Solution:

Ans. (ii), (iii) and (iv)

**BEGINNER'S BOX-6**

- If $\Delta G^\circ > 0$ for a reaction then :-
 (1) $K_p > 1$ (2) $K_p < 1$ (3) $K_p = 1$ (4) None
CTD251
- For an endothermic reaction to be spontaneous :-
 (1) ΔG must be +ve (2) ΔS must be > 0 (3) $T\Delta S$ must be -ve (4) $T\Delta S$ must be equal to ΔG
CTD252
- The value of ΔG for the process $\text{H}_2\text{O}(\text{s}) \longrightarrow \text{H}_2\text{O}(\ell)$ at 1 atm and 260 K is :-
 (1) < 0 (2) $= 0$ (3) > 0 (4) Unpredictable
CTD253
- In a certain chemical reaction $\Delta H = 150 \text{ kJ}$ and $\Delta S = 10 \text{ JK}^{-1}$ at 300 K. The value of ΔG would be :-
 (1) -2850 J (2) Zero (3) $+2850 \text{ J}$ (4) 147 kJ
CTD254
- The standard Gibbs's energy change for a gaseous reaction at 27°C is $X \text{ kCal}$. If equilibrium constant for a reaction is 100 and R is $2 \text{ Cal K}^{-1} \text{ mol}^{-1}$. Then X is :-
 (1) -2.7636 (2) $+2.7636$ (3) $+ 807$ (4) $- 807$
CTD255
- The favourable conditions for a spontaneous reaction are :-
 (1) $T\Delta S > \Delta H$, $\Delta H = +\text{ve}$, $\Delta S = +\text{ve}$ (2) $T\Delta S > \Delta H$, $\Delta H = +\text{ve}$, $\Delta S = -\text{ve}$
 (3) $T\Delta S = \Delta H$, $\Delta H = -\text{ve}$, $\Delta S = -\text{ve}$ (4) $T\Delta S = \Delta H$, $\Delta H = +\text{ve}$, $\Delta S = +\text{ve}$
CTD256

Third Law of Thermodynamics (TLOT) :

At zero kelvin (absolute zero temperature), the entropy of pure perfect crystalline solid is taken as zero.


Exceptions :

(i) NO, N₂O


(ii) CO, CO₂

(iii) Mixture of isotopes

(iv) Ice



Perfect Crystal $\Rightarrow S = 0$



CO Crystal $\Rightarrow S \neq 0$

Introduction to thermochemistry :

Thermochemistry is the branch of physical chemistry which deals with the transfer of heat between a chemical system and its surrounding when a change of phase or chemical reaction takes place within the system.

Depending upon the conditions under which the reaction is carried out, the quantity of heat transferred is related to energy or enthalpy change due to changes of states which occur in the system.


In this chapter we will introduce enthalpies of some specific reactions. Like, Enthalpy of formation (ΔH_f), Enthalpy of combustion (ΔH_{comb}), Bond dissociation enthalpy (ΔH_{BDE}) & Enthalpy of Neutralisation ($\Delta H_{\text{neutralization}}$)

Thermochemical Reaction :

The balanced chemical reaction which give information about the physical states of reactants & products and heat change is called as thermo chemical reaction.



Thermo chemical reactions are of 2 type:




(i) Endothermic reaction :

The reaction which absorbs heat is called as endothermic reaction.

$$\Delta H = +ve$$

$$\Delta H = H_p - H_R = +ve \text{ i.e. } H_p > H_R$$

- Stability of reactant > Stability of product because more heat is required to break the bonds of reactant.
- The product formed in the endothermic reaction is called an endothermic compound.
- If more heat is absorbed then the product formed in the reaction will be less stable or the reactant is more stable.

Representation of endothermic reaction :**Examples :**


(I) Dissociation reactions (mostly)

(II) Fusion reactions

(III) Vaporization reactions

(IV) Sublimation reactions

(V) Photosynthesis

**(ii) Exothermic reaction :**


The reaction in which heat is evolved is called as an exothermic reaction.

$$\Delta H = -ve \quad \Delta H = H_P - H_R = -ve \text{ i.e. } H_P < H_R$$

(a) Stability of reactant < Stability of product because less heat is required to break the bonds of reactant.

(b) The product formed in the exothermic reaction is called an exothermic compound.


(c) If more heat is released then the product formed in the reaction will be more stable or the reactant is less stable.

Representation of exothermic reaction.**Examples :**

(I) Combustion reactions


(II) Neutralisation reactions

(III) Respiration reaction




(IV) Formation $\begin{cases} \nearrow \text{endo} \\ \searrow \text{exo (generally)} \end{cases}$

Exceptions of formation reaction:





★ Golden Key Points ★

- (i) If conditions are not given, then ΔH is considered to be ΔH° .
 (ii) If thermochemical reaction is multiplied by a coefficient then, ΔH of reaction is also multiplied by that coefficient.



- (iii) If reaction is reversed, then numerical value of ΔH remains same but sign is changed.




BEGINNER'S BOX-7

- An endothermic reaction is one in which :
 (1) Heat is converted into electricity (2) Heat is absorbed
 (3) Heat is evolved (4) Heat is converted into mechanical work
CTD257
- If heat of reaction $\text{A} + 5\text{B} \longrightarrow 2\text{C} + 3\text{D}$, is -50 kJ . What is the heat of the reaction $2\text{A} + 10\text{B} \longrightarrow 4\text{C} + 6\text{D}$.
 (1) -50 kJ (2) -25 kJ (3) -100 kJ (4) $+100 \text{ kJ}$
CTD258
- The process $\text{CH}_3\text{COOH} \longrightarrow \text{CH}_3\text{COO}^- + \text{H}^+$, should be :
 (1) Exothermic
 (2) Endothermic
 (3) Neither exothermic nor endothermic
 (4) Exothermic or endothermic depending upon temperature
CTD259
- For the given reaction :
 $\text{CO}_2(\text{g}) + \text{H}_2(\text{g}) \longrightarrow \text{CO}(\text{g}) + \text{H}_2\text{O}(\text{g}) ; \quad \Delta H = 40 \text{ kJ}$
 The ΔH is specifically called
 (1) Heat of formation of CO (2) Heat of combustion
 (3) Heat of reaction (4) Heat of hydrogenation of C = O bond
CTD260

Heat of Reaction or Enthalpy of Reaction (ΔH_R) :

The amount of heat evolved or absorbed when number of moles of the reactant according to the balanced chemical reaction had completely reacted is called as heat of reaction.





Note: Heat of reaction at constant pressure is ΔH and heat of reaction at constant volume is ΔE .

Factors affecting heat of reaction :**(i) Reaction condition :**

The chemical reactions are carried out at constant temperature with either pressure or volume constant. At constant pressure $q_p = \Delta H_{\text{reaction}}$

At constant volume $q_v = \Delta E_{\text{reaction}}$

$$\Delta H = \Delta E + \Delta n_g RT$$

(ii) Quantity of reactant :

Note: If equation is multiplied by a coefficient then value of ΔH is also multiplied by that coefficient.

(iii) Physical state of products and reactants :

If the physical state of product is different then the value of ΔH is different.

Note: For H_2O (liq.), ΔH is more negative in comparison to the formation of H_2O (vap.) because when vapours convert into liquid then some heat is released.

(iv) Allotropic form : (Physical nature of reactant)**(v) Temperature :**

Effect of temperature on heat of reaction is given by **Kirchoff equation**

(i) at constant pressure : $\frac{\Delta H_{T_2} - \Delta H_{T_1}}{T_2 - T_1} = \Delta C_p$

$$\Delta C_p = \Sigma(C_p)_P - \Sigma(C_p)_R$$

ΔH_{T_1} = Heat of reaction at T_1 temperature

ΔH_{T_2} = Heat of reaction at T_2 temperature

(ii) at constant volume : $\frac{\Delta E_{T_2} - \Delta E_{T_1}}{T_2 - T_1} = \Delta C_v$




$$\Delta C_v = \Sigma(C_v)_P - \Sigma(C_v)_R$$

Types of Heat of Reaction :

(A) Heat of formation, Enthalpy of formation (ΔH_f) or ($\Delta_f H$) :

It is the enthalpy change when one mole of a substance is formed from its elements in their most abundant naturally occurring form or in their standard and stable state form (also called reference states).

The reference state of oxygen, carbon and sulphur are O_2 gas, C_{graphite} and S_{rhombic} respectively. Some reactions with standard molar enthalpies of formation are :



APPLICATION OF ΔH_f :

Calculation of ΔH of any general reaction.

Let us consider a general reaction $aA + bB \rightarrow cC + dD$

$$\begin{aligned} \Delta H_{\text{reaction}} &= \sum \Delta H_{f(\text{products})} - \sum \Delta H_{f(\text{reactant})} \\ &= [c\Delta H_{f(C)} + d\Delta H_{f(D)}] - [a\Delta H_{f(A)} + b\Delta H_{f(B)}] \end{aligned}$$

★ Golden Key Points ★

- Standard condition means, $P = 1 \text{ atm}$, $T = 25^\circ\text{C}$ or 298 K
Standard heat of formation is represent by ΔH_f° .
- If no condition is given then value of ΔH_f is considered as ΔH_f° .
- Standard heat of formation of all the elements in stable standard state is taken to be zero.
- The reference state of commonly used elements are

Elements	Reference state
C	$C_{(\text{graphite})}$
S	$S_{8(\text{Rhombic})}$ (Rhombic sulphur is energy wise more stable as compared to monoclinic sulphur)
P	$P_{4(\text{white})}$
O	$O_{2(g)}$
H	$H_{2(g)}$
Br	$Br_{2(l)}$
Metal	$M_{(s)}$ [except $Hg_{(l)}$]

- The formation reaction may be exothermic or endothermic.

Illustration 34:

Since enthalpy of elements in their natural state is taken as zero, the value of ΔH_f of compounds :





- | | |
|---------------------------------|------------------------|
| (1) is always negative | (2) is always positive |
| (3) may be positive or negative | (4) is zero |

Solution:

Ans. (3)

Illustration 35:





The enthalpy of formation of ammonia at 298K is given as $\Delta H_f^\circ = -46.11$ kJ per mol of $\text{NH}_3(\text{g})$. To which of the following equation does this value apply ?

**Solution:**

Ans. (1)

Illustration 36:





Which of the following equation represents the standard heat of formation ?

**Solution:**

Ans. (2)

Illustration 37:

Which of the following reaction defines ΔH_f° ?

**Solution:**

Ans. (2)

Illustration 38:

How much heat will be required at constant pressure to form 1.28 kg of CaC_2 from $\text{CaO}(\text{s})$ & $\text{C}(\text{s})$?

Given :

$$\Delta_f H^\circ (\text{CaO}, \text{s}) = -152 \text{ kCal mol}^{-1}$$

$$\Delta_f H^\circ (\text{CaC}_2, \text{s}) = -14 \text{ kCal mol}^{-1}$$


$$\Delta_f H^\circ (\text{CO}, \text{g}) = -26 \text{ kCal mol}^{-1}$$

$$(1) +112 \text{ kCal}$$

$$(2) 224 \text{ kCal}$$

$$(3) 3840 \text{ kCal}$$

$$(4) 2240 \text{ kCal}$$

Solution:


$$\Delta_f H^\circ = (-14 - 26) - (-152) = +112 \text{ kCal mol}^{-1}$$

$$\text{Total heat required} = \left(\frac{1280}{64} \right) \times 112$$


$$\Rightarrow 2240 \text{ kCal}$$

Illustration 39:

The $\Delta_f H^\circ$ (N_2O_5 , g) in kJ mol^{-1} on the basis of the following data is :



$$\Delta_r H^\circ = -114 \text{ kJ mol}^{-1}$$



$$\Delta_r H^\circ = -102.6 \text{ kJ mol}^{-1}$$

$$\Delta_f H^\circ (\text{NO, g}) = 90.2 \text{ kJ mol}^{-1}$$


(1) 15.1

(2) 30.2


(3) - 36.2

(4) none of these

Solution:




$$\Delta_f H^\circ = 90.2$$




$$\Delta_r H^\circ = 90.2 \times 2$$

... (1)



$$\Delta_r H^\circ = -114$$


... (2)



$$\Delta_r H^\circ = \frac{-102.6}{2} = -51.3$$

... (3)

From Equations (1) + (2) + (3)



$$\Delta_f H^\circ (\text{N}_2\text{O}_5, \text{g}) = 15.1 \text{ kJ mol}^{-1}$$

Illustration 40:

Calculate ΔH° for $2\text{Al}_{(s)} + \text{Fe}_2\text{O}_3 \longrightarrow 2\text{Fe}_{(s)} + \text{Al}_2\text{O}_3$ given that standard enthalpy of Fe_2O_3 and Al_2O_3 are -196.5 and -399.1 kCal.

Solution:

$$\Delta H_{\text{Reaction}}^\circ = \Sigma \Delta H_{\text{P}}^\circ - \Sigma \Delta H_{\text{R}}^\circ$$


$$= [2 \times \Delta H_{\text{Fe(s)}}^\circ + \Delta H_{\text{Al}_2\text{O}_3}^\circ] - [2 \times \Delta H_{\text{Al(s)}}^\circ + \Delta H_{\text{Fe}_2\text{O}_3}^\circ]$$

$$= 2 \times 0 + (-399.1) - [2 \times 0 + (-196.5)]$$

$$\Delta H_{\text{Reaction}}^\circ = -202.6 \text{ kCal}$$

Illustration 41:

The heat of formation of the compound in the following reaction is :



(1) -44 kCal mol^{-1}

(2) -22 kCal mol^{-1}

(3) +11 kCal mol^{-1}

(4) -88 kCal mol^{-1}

Solution:





Ans. (2)

For the formation of 1 mol of HCl from elements $\Delta H_f^\circ = \frac{-44}{2} = -22 \text{ kCal}$

Heat of combustion (ΔH_{comb}) :

Amount of heat evolved when 1 mole of substance is completely burnt (or oxidised) in excess of oxygen.

Example :



Note :

- (I) Heat of combustion reaction is always exothermic
- (II) If conditions are not given then ΔH_{comb} considered as $\Delta H^{\circ}_{\text{comb}}$.
- (III) If in a reaction heat of combustion of reactants and products are given then heat of that reaction can be measured as follows

$$\Delta H = \sum (\Delta H_{\text{comb}})_{\text{R}} - (\Delta H_{\text{comb}})_{\text{P}}$$

Application of Heat of Combustion :**Calorific value or fuel value (C.V.) :**

The amount of heat evolved when 1 g of a substance (food or fuel) is completely burnt (or oxidised)

$$\text{Calorific value} = \frac{\Delta H_{\text{comb}}}{\text{Molecular weight}}$$

Unit :- kJ g⁻¹ or kCal g⁻¹

★ **Golden Key Points** ★

- Heat of combustion reaction is always exothermic
- If conditions are not given then ΔH_{comb} is considered as $\Delta H^{\circ}_{\text{comb}}$.
- Maximum value of calorific value = Maximum efficiency or best fuel
- H₂ has the highest calorific value (150 kJ/g) but it is not used as domestic or industrial fuel due to some technical problems.


Illustration 42:

Calculate the heat of formation of Benzene. The reaction is given as $6\text{C}(\text{s}) + 3\text{H}_2(\text{g}) \rightarrow \text{C}_6\text{H}_6(\ell)$ and -3268 , -393.5 and -285.8 kJ mol⁻¹ are the heats of combustion of benzene, heat of formation of CO₂ and heat of formation of H₂O(ℓ) respectively.

Solution:

Target reaction, $6\text{C}(\text{s}) + 3\text{H}_2(\text{g}) \rightarrow \text{C}_6\text{H}_6(\ell)$

Given



$6 \times \text{eq. (ii)} + 3 \times \text{eq. (iii)} - \text{eq. (i)}$



$$\Delta H = 6 \times -393.5 + 3 \times -285.8 - (-3268) = +49.6 \text{ kJ mol}^{-1}$$

Illustration 43:

The heats of formation of $\text{CO}_2(\text{g})$ and $\text{H}_2\text{O}(\ell)$ are -97 and $-68 \text{ kCal mol}^{-1}$. The heat of combustion of benzene is $-783 \text{ kCal mol}^{-1}$. What will be the heat of formation of benzene?

Solution:

Given :



Target reaction



$$6 \times \text{eq. (ii)} + 3 \times \text{eq. (iii)} - \text{eq. (i)}$$



$$\Delta H = 6 \times -97 + 3 \times -68 - (-783) = -3 \text{ kCal mol}^{-1}$$

Illustration 44:


Calculate the enthalpy of combustion of ethylene (gas) to form CO_2 (gas) and H_2O (gas) at 298 K and 1 atmospheric pressure. The enthalpies of formation of $\text{CO}_2(\text{g})$, $\text{H}_2\text{O}(\text{g})$ and $\text{C}_2\text{H}_4(\text{g})$ are -393.7 , -241.8 , $+52.3 \text{ kJ}$ per mol respectively.

Solution:

We are given :



Target reaction



$2 \times \text{Equation (i)} + 2 \times \text{Equation (ii)} - \text{Equation (iii)}$ gives

$$\Delta H = 2(-393.7) + 2(-241.8) - (52.3) = -1323.3 \text{ kJ mol}^{-1}$$


Illustration 45:

1 mole of methanol, when burnt in oxygen, gives out -723 kJ mol^{-1} heat. If 1 mole of oxygen is used what will be the amount of heat evolved?

- (1) 723 kJ (2) 964 kJ (3) 482 kJ (4) 241 kJ

Solution:

Ans. (3)



$$\text{with } 1 \text{ mole of O}_2, \quad \Delta H = \frac{-723 \times 2}{3} = -482 \text{ kJ}$$

Illustration 46:

Combustion of methane :

(1) is an exothermic reaction

(2) is an endothermic reaction

(3) requires a catalyst

(4) gives H_2


Solution:

Ans. (1)

Combustion is always exothermic.

Illustration 47:

The heat evolved in the combustion of glucose is given by the equation



The weight of $CO_2(g)$ produced when 170 kCal of heat is evolved in the combustion of glucose is

(1) 264 g

(2) 66 g

(3) 11 g

(4) 44 g

Solution:


Ans. (2)

Evolution of 680 kCal is accompanied by $CO_2 = 6 \times 44 = 264 \text{ g}$

Evolution of 170 kCal is accompanied by $CO_2 = \frac{264}{680} \times 170 = 66 \text{ g}$

Illustration 48:

Find out the calorific value of Glucose

**Solution:**

$\therefore$ Heat evolved from 1 mol glucose = 2900 kJ

or Heat evolved from 180 gram glucose = 2900 kJ

$\therefore$ Heat evolved from 1 gram glucose = $\frac{2900}{180} = 16.11 \text{ kJ g}^{-1}$

or another method $C.V. = \frac{\Delta H_{\text{comb}}}{M_w} = \frac{2900}{180} = 16.11 \text{ kJ g}^{-1}$

Illustration 49:

Enthalpy of combustion of a substance is always :

(1) > 0

(2) > 0

(3) < 0

(4) < 0

Solution:

Ans. (4)

Illustration 50:

The heat change for a reaction : $CO(g) + \frac{1}{2} O_2 \longrightarrow CO_2(g)$ refers to

(1) enthalpy of formation of carbon dioxide

(2) enthalpy of combustion of carbon dioxide

(3) enthalpy of vapourisation



(4) enthalpy of combustion of carbon monoxide

Solution:

Ans. (4)


Heat of Neutralization (ΔH_{neut}) :

The heat evolved when one equivalent of an acid is completely neutralized by one equivalent of a base in dilute solution is called as heat of neutralization.




Note :

- (i) When one equivalent of SA is neutralized by one equivalent of SB, then heat evolved remains constant and its value is $-13.7 \text{ kCal/equivalent}$ or $-57.2 \text{ kJ equivalent}^{-1}$.



- (ii) If one of the acid or base or both are weak, then heat of neutralization is usually less than $-13.7 \text{ kCal eq}^{-1}$ or -57.3 kJ eq^{-1} because some part of the heat released in neutralization is absorbed to dissociate the weak electrolyte completely.



Exception : For a reaction $\text{HF} + \text{NaOH} \rightarrow \text{NaF} + \text{H}_2\text{O}; \Delta H = -16.7 \text{ kCal}$; this is because of hydration of F^- ion.

Illustration 51:

Heat of neutralisation of an acid by a base is maximum when :

- (1) Both the acid and base are weak
- (2) Both the acid and base are strong
- (3) The acid is strong and the base is weak
- (4) The acid is weak and the base is strong

Solution:

Ans. (2)

Heat of neutralisation is maximum (57.2 kJ eq^{-1} or $13.7 \text{ kCal eq}^{-1}$) when both acid and base are strong.

Illustration 52:

If $\text{H}^+ + \text{OH}^- \longrightarrow \text{H}_2\text{O} + 13.7 \text{ kCal}$, then heat of complete neutralisation of 1 gm mol of H_2SO_4 with base in excess will be :

- (1) -13.7 kCal (2) -27.4 kCal (3) -6.85 kCal (4) -3.425 kCal

Solution:

Ans. (2)


Moles of $\text{H}_2\text{SO}_4 = 1 \text{ mol}$

g eq. of $\text{H}_2\text{SO}_4 = \text{moles} \times \text{V.F.} = 1 \times 2 = 2 \text{ g eq.}$

Heat evolve due to 2 g eq. = $-13.7 \times 2 = -27.4 \text{ kCal}$

Illustration 53:

200 cm³ of 0.1 M H₂SO₄ is mixed with 150 cm³ of 0.2 M KOH. Find the value of evolved heat.

Solution:

Heat liberated by 1 eq. = 57.2 kJ

So heat liberated by 0.03 eq. = 57.2 × 0.03 = 1.7 kJ

Heat of Hydrogenation ($\Delta H_{\text{Hydrogenation}}$) :

The heat evolved during the complete hydrogenation of one mol unsaturated organic compound into its saturated compound is called as heat of hydrogenation.

Unsaturated organic compound $\xrightarrow{\text{Change}}$ Saturated organic compound

(= or $\equiv$ Bond)


(– Bond)



Note : Heat of hydrogenation is exothermic process.

Heat of Atomization (ΔH_{atom}) :


The amount of heat required to dissociate 1 mol substance into gaseous atoms is called as heat of atomization.



Note : It is an endothermic reaction.


Heat of Transformation :

- (i) **Heat of fusion (ΔH_{fusion}) :** The required amount of heat to convert 1 mole solid into liquid at its melting point is called as heat of fusion.



Note : Heat of fusion is always endothermic reaction i.e. ($\Delta H = +ve$)


- (ii) **Heat of vaporization (ΔH_{vapour}) :** The required amount of heat to convert 1 mole liquid into gas at its boiling point is called as heat of vaporization.



Note : Heat of vaporization is always endothermic reaction i.e. ($\Delta H = +ve$)

(iii) Heat of sublimation (ΔH_{Sub}) :

The required amount of heat to convert 1 mole solid into gas at a certain temperature is called as heat of sublimation.



Note : Heat of sublimation is always endothermic reaction i.e. ($\Delta H = + \text{ve}$)

Illustration 54:

The enthalpy change for the process $\text{C}_{(\text{s})} \longrightarrow \text{C}_{(\text{g})}$ corresponds to the enthalpy of

- (1) fusion (2) vapourization (3) combustion (4) sublimation

Solution:

Ans. (4)


Solid $\longrightarrow$ gas, is sublimation.

**BEGINNER'S BOX-8**

- Enthalpy of neutralisation of acetic acid with KOH will be numerically :
(1) = 57.2 kJ (2) > 57.2 kJ (3) < 57.2 kJ (4) unpredictable CTD261
- The vapourisation process is always :
(1) Exothermic (2) Endothermic
(3) Can be exothermic or endothermic (4) None of these CTD262
- One mole of H_2SO_4 is completely neutralised with 2 mole of NaOH in dilute solutions. The amount of heat evolved during the process is :
(1) 57.2 kJ (2) $\frac{57.2\text{kJ}}{2}$ (3) 13.7 kCal (4) 114.4 kJ CTD263
- Which of the following data represents the value of heat of neutralisation of strong acid against strong base ?
(1) - 13.7 kCal (2) - 57.2 kJ (3) - 5.72×10^4 J (4) All the above CTD264
- Fusion of ice is :
(1) Exothermic change
(2) Endothermic change
(3) A process that does not involve any heat change
(4) Unpredictable CTD265

Bond Energy / Bond Dissociation Energy :

The required amount of energy to dissociate **one mole gaseous bond** into separate **gaseous atoms** is called as **bond dissociation energy**.





★ Golden Key Points ★

- The **bond energy** may be defined as the **average** amount of energy required to dissociate one mole gaseous bond into separate gaseous atoms.
- Bond dissociation process is an endothermic process.
- If bond energy of various bonds present in the reactants and products are given then ΔH of that reaction can be calculate as follows.

$$\Delta H = \Sigma(\text{B. E.})_R - \Sigma(\text{B. E.})_P$$

- In the case of poly atomic molecule** we calculate the average bond energy.

$$(\text{BE})_{\text{av}} = \text{Average bond energy} = \frac{\text{Total energy required with all bonds}}{\text{Number of bond dissociation}}$$



$$(\text{BE})_{\text{av}} = \text{Average bond energy} = \frac{112+108}{2} = 110 \text{ kJ mol}^{-1}$$

Illustration 55:

Given the bond energy of $\text{N}\equiv\text{N}$, $\text{H}-\text{H}$ and $\text{N}-\text{H}$ bonds are 945, 436 and 391 kJ mol^{-1} respectively, the enthalpy of the reaction $\text{N}_{2(\text{g})} + 3\text{H}_{2(\text{g})} \longrightarrow 2\text{NH}_{3(\text{g})}$ is :

(1) - 93 kJ


(2) 102 kJ

(3) 90 kJ

(4) 105 kJ

Solution:

Ans. (1)





$$945 + 3 \times 436 \qquad 2 \times (3 \times 391)$$

$$= 2253 \text{ kJ} \qquad = 2346 \text{ kJ}$$

$$\Delta H = \Sigma (\text{B.E.})_R - \Sigma (\text{B.E.})_P = 2253 - 2346 = -93 \text{ kJ}$$

Illustration 56:

The enthalpy changes at 298 K in successive breaking of O-H bonds of $\text{H}-\text{O}-\text{H}$ are



The bond enthalpy of the O - H bond is

(1) 498 kJ mol^{-1}

(2) 463 kJ mol^{-1}

(3) 428 kJ mol^{-1}

(4) 70 kJ mol^{-1}



Solution:

Ans. (2)

$$(\text{B.E.})_{\text{av}} = \frac{498 + 428}{2} = 463 \text{ kJ}$$

Illustration 57:

The required heat for dissociation of 1 mol H_2O into its atoms (H and oxygen) is $\Delta H_{\text{Dis.}}$. Then calculate the bond energy of O-H bond.



**Solution:**

$\therefore$ required energy for breaking the 2 mol O - H bond = $\Delta H_{\text{Dis.}}$

$$\therefore \text{required energy for 1 mole} = \frac{\Delta H_{\text{Dis.}}}{2}; \Delta H_{\text{O-H}} = \frac{\Delta H_{\text{Dis.}}}{2}$$

Illustration 58:

Calculate the bond energy of C-H Bond in methane.

Solution:

$\therefore$ Bond energy of 4 mol C-H = ΔH_{Dis}

$$\therefore \text{Bond energy of 1 mol C-H bond} = \frac{\Delta H_{\text{Dis.}}}{4}$$

Illustration 59:

The energy change of reaction $\text{C}_2\text{H}_{6(\text{g})} \longrightarrow 2\text{C}_{(\text{g})} + 6\text{H}_{(\text{g})}$ is X kJ. The bond energy of C-H bond is:

- (1) $\frac{X}{6} \text{ kJ mol}^{-1}$ (2) $\frac{X}{3} \text{ kJ mol}^{-1}$ (3) $X \text{ kJ /mol}^{-1}$ (4) unpredictable from data

Solution:

Ans. (4)

If we know C-C bond energy than only we can calculate C-H bond energy using given data.

Illustration 60:

The value of ΔH represents :

- (1) enthalpy of solution of copper (II) sulphate
- (2) enthalpy of hydration of copper (II) sulphate
- (3) enthalpy of hydrolysis of copper (II) sulphate
- (4) lattice energy of copper (II) sulphate

Solution:

Ans. (2)

Illustration 61:

The bond energy of hydrogen is 103 kCal mol⁻¹. This means that :

- (1) 103 kCal are required to break 6.023×10^{23} gaseous H_2 molecules into gaseous atoms
- (2) 103 kCal are required to break the bonds in one gram of hydrogen
- (3) 103 kCal are required to break one bond to form two atoms of hydrogen
- (4) 103 kCal are required to break one mole of gaseous hydrogen molecules into ions.

Solution:

Ans. (1)

Heat of Hydration (ΔH_{hydra}) :

Amount of heat evolved when **one mole** of anhydrous salt combines with fixed number of water molecules to convert into its specific hydrated crystal is called as heat of hydration.

Example :

- $1 \text{ CuSO}_4(\text{s}) + 5\text{H}_2\text{O}(\ell) \longrightarrow \text{CuSO}_4 \cdot 5 \text{ H}_2\text{O}(\text{s})$ $\Delta H = -ve$
 anhydrous salt hydrated salt
- $1 \text{ MgSO}_4(\text{s}) + 7\text{H}_2\text{O} (\ell) \longrightarrow \text{MgSO}_4 \cdot 7\text{H}_2\text{O}(\text{s})$ $\Delta H = -ve$
 anhydrous salt hydrated salt
- $1 \text{ CaCl}_2(\text{s}) + 6\text{H}_2\text{O}(\ell) \longrightarrow \text{CaCl}_2 \cdot 6 \text{ H}_2\text{O}(\text{s})$ $\Delta H = -ve$
 anhydrous salt hydrated salt

Special Note : Heat of hydration is exothermic

Heat of Solution ($\Delta H_{\text{sol.}}$) :



Amount of heat absorbed or evolved when **one mol** of substance is dissolved in such a large volume of solvent that further addition of solvent does not produce any more heat change is called as '**Heat of solution**'.

Example 1:	$1 \text{ CuSO}_{4(s)} + \text{aq}$	$\longrightarrow$	$\text{CuSO}_4(\text{aq})$	$\Delta H_{\text{solution}}$	✓
Example 2:	$\text{MgSO}_{4(s)} + 7\text{H}_2\text{O}(\ell) + \text{aq}$	$\longrightarrow$	$\text{MgSO}_4(\text{aq})$	$\Delta H_{\text{solution}}$	✓
	$\text{MgSO}_4 \cdot 7\text{H}_2\text{O}(s) + \text{aq}$	$\longrightarrow$	$\text{MgSO}_4(\text{aq})$	$\Delta H_{\text{solution}}$	✓


Special Note : Heat of solution may be endothermic or exothermic.

Illustration 62:

The heat of solution of anhydrous $\text{CuSO}_4(\text{s})$ is $-15.9 \text{ kCal mol}^{-1}$ and that of $\text{CuSO}_4 \cdot 5\text{H}_2\text{O}(\text{s})$ is $2.8 \text{ kCal mol}^{-1}$. Calculate the heat of hydration of $\text{CuSO}_4(\text{s})$.

Solution:

Subtracting Eq. (ii) from Eq.(i)




Heat of hydration of $\text{CuSO}_4 = -18.7 \text{ kCal mol}^{-1}$

Laws of Thermochemistry :


(I) Lavoisier and Laplace Law :

Enthalpy of formation of compound is **numerically equal** to the enthalpy of decomposition of that compound with **opposite sign**.



(II) Hess Law of Constant Heat Summation :

The heat change in a complete chemical reaction always remain same whether reaction completes in one step or more.




$$\Delta H = \Delta H_1 + \Delta H_2$$

or
$$\Delta H = \Delta H_3 + \Delta H_4 + \Delta H_5$$

or
$$\Delta H = \Delta H_1 + \Delta H_2 = \Delta H_3 + \Delta H_4 + \Delta H_5$$

Example:




★ Golden Key Points ★

- Heat change of a reaction does not depend on the **number of steps** used in the reaction.
- Heat change of a reaction does not depend on **intermediate position**, it depend only on initial and final state.
- Heat change of a chemical reaction does not depend on **time of reaction**.

Illustration 63:

Single step reaction $A \rightarrow B$; $\Delta H = ?$

Multi step reaction to produce B from A is given



Solution:

According to Hess's law $\Delta H = \Delta H_1 + \Delta H_2 + \Delta H_3 + \Delta H_4$



BEGINNER'S BOX-9

1. A hypothetical reaction, $X \longrightarrow 2Y$ proceeds by the following sequence of steps





The value of ΔH of reaction is :

- (1) $q_1 + q_2 + q_3$ (2) $2q_1 + 2q_2 + 3q_3$ (3) $2(q_1 + q_2 + 2q_3)$ (4) $2(q_1 + q_2 + q_3)$

CTD266

2. Consider two paths of a certain reaction



(1) $(\Delta H_1 + \Delta H_2) > (\Delta H_3 + \Delta H_4)$

(2) $(\Delta H_1 + \Delta H_2) = (\Delta H_3 + \Delta H_4)$

(3) $(\Delta H_2 + \Delta H_3) = (\Delta H_1 + \Delta H_4)$

(4) $(\Delta H_1 + \Delta H_2) < (\Delta H_3 + \Delta H_4)$

CTD267



BEGINNER'S BOX

ANSWERS KEY

BEGINNER'S BOX-1	Que.	1	2	3	4					
	Ans.	1	1	1	1					
BEGINNER'S BOX-2	Que.	1	2	3	4					
	Ans.	3	4	2	1					
BEGINNER'S BOX-3	Que.	1	2	3	4	5	6	7	8	
	Ans.	2	2	2	3	1	2	4	3	
BEGINNER'S BOX-4	Que.	1	2							
	Ans.	1	3							
BEGINNER'S BOX-5	Que.	1	2	3	4	5	6	7		
	Ans.	2	3	4	3	3	1	2		
BEGINNER'S BOX-6	Que.	1	2	3	4	5	6			
	Ans.	2	2	3	4	1	1			
BEGINNER'S BOX-7	Que.	1	2	3	4					
	Ans.	2	3	2	3					
BEGINNER'S BOX-8	Que.	1	2	3	4	5				
	Ans.	3	2	4	4	2				
BEGINNER'S BOX-9	Que.	1	2							
	Ans.	3	2							