

Exercise - I (Conceptual Questions)

Build Up Your Understanding

CALCULATION OF CENTRE OF MASS

1. Three identical spheres, each of mass 1 kg are placed touching each other with their centres on a straight line. Their centre are marked P, Q and R respectively. The distance of centre of mass of the system from P is :

(1) $\frac{PQ + PR + QR}{3}$ (2) $\frac{PQ + PR}{3}$
 (3) $\frac{PQ + QR}{3}$ (4) $\frac{PR + QR}{2}$


PSC001

2. A uniform metal disc of radius R is taken and out of it a disc of diameter $\frac{R}{2}$ is cut off from the end. The centre of mass of the remaining part will be:

(1) $\frac{R}{10}$ from the centre
 (2) $\frac{R}{15}$ from the centre
 (3) $\frac{R}{5}$ from the centre
 (4) $\frac{R}{20}$ from the centre

PSC002

3. The coordinate of the centre of mass of a system as shown in figure :-



(1) $\left(\frac{a}{3}, 0\right)$ (2) $\left(\frac{a}{2}, \frac{a}{2}\right)$
 (3) $\left(\frac{a}{3}, \frac{a}{3}\right)$ (4) $\left(0, \frac{a}{3}\right)$

PSC003

4. The centre of mass of a system of particles does not depend on :

- (1) masses of the particles
 (2) Internal forces on the particles
 (3) position of the particles
 (4) relative distance between the particles

PSC004

5. The centre of mass of a system of two particles divides the distance between them

- (1) In inverse ratio of square of masses of particles
 (2) In direct ratio of square of masses of particles
 (3) In inverse ratio of masses of particles
 (4) In direct ratio of masses of particles

PSC005

6. The centre of mass of a body :-

- (1) Lies always outside the body
 (2) May lie within, outside of the surface of the body
 (3) Lies always inside the body
 (4) Lies always on the surface of the body

PSC006

7. Three identical metal balls, each of radius r, are placed touching each other on a horizontal surface such that an equilateral triangle is formed when the centres of the three balls are joined. The centre of mass of the system is located at :-

- (1) horizontal surface
 (2) centre of one of the balls
 (3) line joining centres of any two balls
 (4) point of intersection of their medians

PSC007

8. A system consists of mass M and m ($m \ll M$). The centre of mass of the system is :-

- (1) at the middle
 (2) nearer to M
 (3) nearer to m
 (4) at the position of larger mass.

PSC008

9. The centre of mass of a system of three particles of masses 1g, 2g and 3g is taken as the origin of a coordinate system. The position vector of a fourth particle of mass 4g such that the centre of mass of the four particle system lies at the point $(1, 2, 3)$ is $\alpha(\hat{i} + 2\hat{j} + 3\hat{k})$, where α is a constant. The value of α is :-
- (1) $\frac{10}{3}$ (2) $\frac{5}{2}$ (3) $\frac{1}{2}$ (4) $\frac{2}{5}$

PSC009

MOTION OF CENTRE OF MASS

10. The law of conservation of momentum for a system is based on Newton's :-
- (1) First law of motion
(2) Second law of motion
(3) Third law of motion
(4) Law of gravitation

PSC010

11. A person of mass m is standing on one end of a plank of mass M and length L and floating in water. The person moves from one end to another and stops. The displacement of the plank is -
- (1) $\frac{Lm}{(m+M)}$ (2) $Lm(M+m)$
(3) $\frac{(M+m)}{Lm}$ (4) $\frac{LM}{(m+M)}$


PSC011

12. Bullets of mass 40 g each are fired from a machine gun with a velocity of 10^3 m/s. If the person firing the bullets experience an average force of 200N, then the number of bullets fired per minute will be-
- (1) 300 (2) 600 (3) 150 (4) 75

PSC012

13. If the system is released, then the acceleration of the centre of mass of the system :-

- (1) $\frac{g}{4}$
(2) $\frac{g}{2}$
(3) g
(4) $2g$



PSC013

14. Initially two stable particles x and y start moving towards each other under mutual attraction. If at one time the velocities of x and y are V and $2V$ respectively, what will be the velocity of centre of mass of the system?
- (1) V (2) Zero
(3) $\frac{V}{3}$ (4) $\frac{V}{5}$

PSC014

15. A 2 kg body and a 3 kg body are moving along the x -axis. At a particular instant the 2 kg body has a velocity of 3 m/s and the 3 kg body has the velocity of 2 m/s. The velocity of the centre of mass at that instant is :-
- (1) 5 m/s (2) 1 m/s
(3) 0 (4) $\frac{12}{5}$ m/s

PSC015

16. Two objects of masses 200 gram and 500 gram possess velocities $10\hat{i}$ m/s and $3\hat{i} + 5\hat{j}$ m/s respectively. The velocity of their centre of mass in m/s is :-
- (1) $5\hat{i} - 25\hat{j}$ (2) $\frac{5}{7}\hat{i} - 25\hat{j}$
(3) $5\hat{i} + \frac{25}{7}\hat{j}$ (4) $25\hat{i} - \frac{5}{7}\hat{j}$

PSC016

MOMENTUM CONSERVATION

17. A bomb of mass 9 kg explodes into two pieces of 3kg and 6 kg. The velocity of 3 kg piece is 16 m/s. The kinetic energy of 6 kg piece is :-
- (1) 768 J (2) 786 J
(3) 192 J (4) 687 J

PSC017

18. A bomb initially at rest explodes by it self into three equal mass fragments. The velocities of two fragments are $(3\hat{i} + 2\hat{j})$ m/s and $(-\hat{i} - 4\hat{j})$ m/s. The velocity of the third fragment is (in m/s) :-
- (1) $2\hat{i} + 2\hat{j}$ (2) $2\hat{i} - 2\hat{j}$
(3) $-2\hat{i} + 2\hat{j}$ (4) $-2\hat{i} - 2\hat{j}$

PSC018

19. A bomb of 50 Kg is fired from a cannon with a velocity 600 m/s. If the mass of the cannon is 10^3 kg, then its velocity will be –

(1) 30 m/s (2) –30 m/s
(3) 0.30 m/s (4) –0.30 m/s

PSC019

20. A metal ball does not rebound when struck on a wall, whereas a rubber ball of same mass when thrown with the same velocity on the wall rebounds. From this it is inferred that –

(1) Change in momentum is same in both
(2) Change in momentum in rubber ball is more
(3) Change in momentum in metal ball is more
(4) Initial momentum of metal ball is more than that of rubber ball

PSC020

21. A bomb of mass $m = 1$ kg thrown vertically upwards with a speed $u = 100$ m/s explodes into two parts after $t = 5$ s. A fragment of mass $m_1 = 400$ g moves downwards with a speed $v_1 = 25$ m/s, then speed v_2 and direction of another mass m_2 will be :-

(1) 40 m/s downwards (2) 40 m/s upwards
(3) 60 m/s upwards (4) 100 m/s upwards

PSC021

22. A 1 kg stationary bomb is exploded in three parts having mass ratio 1 : 1 : 3. Parts having same mass move in perpendicular directions with velocity 30 m/s, then the velocity of bigger part will be :-

(1) $10\sqrt{2}$ m/s (2) $\frac{10}{\sqrt{2}}$ m/s
(3) $15\sqrt{2}$ m/s (4) $\frac{15}{\sqrt{2}}$ m/s

PSC022

23. A heavy nucleus at rest breaks into two fragments which fly off with velocities 8 : 1. The ratio of radii of the fragments is :-

(1) 1 : 2 (2) 1 : 4 (3) 4 : 1 (4) 2 : 1

PSC023

24. A stationary particle explodes into two particles of masses m_1 and m_2 which move in opposite directions with velocities v_1 and v_2 . The ratio of their kinetic energies E_1/E_2 is :-

(1) m_2/m_1 (2) m_1/m_2
(3) 1 (4) m_1v_2/m_2v_1

PSC024

25. A body of mass 4 m at rest explodes into three pieces. Two of the pieces each of mass m move with a speed v each in mutually perpendicular directions. The total kinetic energy released is :-

(1) $\frac{1}{2}mv^2$ (2) mv^2
(3) $\frac{3}{2}mv^2$ (4) $\frac{5}{2}mv^2$

PSC025

26. A bomb of mass 3.0 kg explodes in air into two pieces of masses 2.0 kg and 1.0 kg. The smaller mass goes at a speed of 80 m/s. The total energy imparted to the two fragments is -

(1) 1.07 kJ (2) 2.14 kJ
(3) 2.4 kJ (4) 4.8 kJ

PSC026

27. A bomb of mass 30 kg at rest explodes into two pieces of masses 18 kg and 12 kg. The velocity of 18 kg mass is 6 m/s. The kinetic energy of the other mass is :-

(1) 524 J (2) 256 J (3) 486 J (4) 324 J

PSC027

28. A bullet of mass m is fired from a gun of mass M . The recoiling gun compresses a spring of force constant k by a distance d . Then the velocity of the bullet is :-

(1) $kd\sqrt{M/m}$ (2) $\frac{d}{M}\sqrt{km}$
(3) $\frac{d}{m}\sqrt{kM}$ (4) $\frac{kM}{m}\sqrt{d}$

PSC028

29. Identify the wrong statement.

(1) A body can have momentum without mechanical energy
(2) A body can have energy without momentum
(3) The momentum is conserved in an elastic collision only.
(4) Kinetic energy is not conserved in an inelastic collision

PSC029

COLLISION

30. A ball strikes the floor and after collision rebounds back. In this state –

(1) Momentum of the ball is conserved
(2) Mechanical energy of the ball is conserved
(3) Momentum of ball–earth system is conserved
(4) The kinetic energy of ball–earth system is conserved

PSC030

31. A bullet of mass P is fired with velocity Q in a large body of mass R . The final velocity of the system will be :-

(1) $\frac{R}{P+R}$ (2) $\frac{PQ}{P+R}$
 (3) $\frac{(P+Q)}{R}$ (4) $\frac{(P+R)}{P}Q$

PSC031

32. A sphere of mass m moving with a constant velocity collides with another stationary sphere of same mass. The ratio of velocities of two spheres after collision will be, if the co-efficient of restitution is e -

(1) $\frac{1-e}{1+e}$ (2) $\frac{e-1}{e+1}$
 (3) $\frac{1+e}{1-e}$ (4) $\frac{e+1}{e-1}$

PSC032

33. Two elastic bodies P and Q having equal masses are moving along the same line with velocities of 16 m/s and 10 m/s respectively. Their velocities after the elastic collision will be in m/s :-

(1) 0 and 25 (2) 5 and 20
 (3) 10 and 16 (4) 20 and 5

PSC033

34. The unit of the coefficient of restitution is -

(1) m/s
 (2) s/m
 (3) $m \times s$
 (4) None of the above

PSC034

35. Two solid balls of rubber A and B whose masses are 200 gm and 400 gm respectively, are moving in mutually opposite directions. If the velocity of ball A is 0.3 m/s and both the balls come to rest after collision, then the velocity of ball B is -

(1) 0.15 m/s
 (2) -0.15 m/s
 (3) 1.5 m/s
 (4) None of the above

PSC035

36. A 1 Kg ball falls from a height of 25 cm and rebounds upto a height of 9 cm. The coefficient of restitution is -

(1) 0.6 (2) 0.32 (3) 0.40 (4) 0.56


PSC036

37. A 50 gm bullet moving with a velocity of 10 m/s gets embedded into a 950 gm stationary body. The loss in kinetic energy of the system will be -

(1) 5% (2) 50% (3) 100% (4) 95%

PSC037

38. A bullet of mass m moving with a speed v strikes a wooden block of mass M and gets embedded into the block. The final speed is :-



(1) $\sqrt{\frac{M}{M+m}}v$ (2) $\sqrt{\frac{m}{M+m}}v$
 (3) $\frac{m}{M+m}v$ (4) $\frac{v}{2}$

PSC038

39. A ball is dropped from height h on the ground level. If the coefficient of restitution is e then the height upto which the ball will go after n^{th} jump will be -

(1) $\frac{h}{e^{2n}}$ (2) $\frac{e^{2n}}{h}$
 (3) he^n (4) he^{2n}

PSC039

40. Two bodies of same mass are moving with same speed V in mutually opposite directions. They collide and stick together. The resultant velocity of the system will be -

(1) Zero (2) $\frac{V}{2}$
 (3) V (4) From Zero to ∞

PSC040

41. The bob (mass m) of a simple pendulum of length L is held horizontal and then released. It collides elastically with a block of equal mass lying on a frictionless table. The kinetic energy of the block will be :-

(1) Zero (2) mgL
 (3) $2mgL$ (4) $\frac{mgL}{2}$

PSC041

42. Two particles each of mass m travelling with velocities u_1 and u_2 collide perfectly inelastically. The loss of kinetic energy will be –

(1) $\frac{1}{2} m(u_1 - u_2)^2$ (2) $\frac{1}{4} m(u_1 - u_2)^2$
 (3) $m(u_1 - u_2)^2$ (4) $2m(u_1 - u_2)^2$

PSC042

43. A ball moving with velocity of 9 m/s collides with another similar stationary ball. After the collision both the balls move in directions making an angle of 30° with the initial direction. After the collision their speed will be –

(1) 2.6 m/s (2) 5.2 m/s
 (3) 0.52 m/s (4) 52 m/s

PSC043

44. A solid sphere is moving and it makes an elastic collision with another stationary sphere of half of its own radius. After collision it comes to rest. The ratio of the densities of materials of second sphere and first sphere is –
- (1) 2 (2) 4 (3) 8 (4) 16

PSC044

45. A 5 kg body collides with another stationary body. After the collision, the bodies move in the same direction with one-third of the velocity of the first body. The mass of the second body will be –

(1) 5 kg (2) 10 kg
 (3) 15 kg (4) 20 kg

PSC045

46. A 10 g bullet, moving with a velocity of 500 m/s, enters a stationary piece of ice of mass 10 kg and stops. If the piece of ice is lying on a frictionless plane, then its velocity will be

(1) 5 cm/s (2) 5 m/s
 (3) 0.5 m/s (4) 0.5 cm/s

PSC046

47. A heavy body moving with a velocity 20 m/s and another small object at rest undergo an elastic collision. The latter will move with a velocity of :-

(1) 20 m/s. (2) 40 m/s.
 (3) 60 m/s. (4) Zero

PSC047

48. A 5 gm lump of clay, moving with a velocity of 10 cm/s towards east, collides head-on with another 2 gm lump of clay moving with 15 cm/s towards west. After collision, the two lumps stick together. The velocity of the compound lump will be –

(1) 5 cm/s towards east
 (2) 5 cm/s towards west
 (3) 2.88 cm/s towards east
 (4) 2.5 cm/s towards west

PSC048

49. In an inelastic collision between two bodies, the physical quantity that is conserved :-

(1) Kinetic energy
 (2) Momentum
 (3) Potential energy
 (4) Kinetic energy and momentum

PSC049

50. A mass of 20 kg moving with a speed of 10 m/s collides with another stationary mass of 5 kg. As a result of the collision, the two masses stick together. The kinetic energy of the composite mass will be :-

(1) 600 J (2) 800 J
 (3) 1000 J (4) 1200 J

PSC050

51. A body of mass m having an initial velocity v makes head on elastic collision with a stationary body of mass M . After the collision, the body of mass m comes to rest and only the body having mass M moves. This will happen only when :-

(1) $m \gg M$ (2) $m \ll M$
 (3) $m = M$ (4) $m = \frac{M}{2}$

PSC051

52. A body A experiences perfectly elastic collision with a stationary body B. If after collision the bodies fly apart in the opposite direction with equal speeds, the mass ratio of A and B is :-

(1) $\frac{1}{2}$ (2) $\frac{1}{3}$ (3) $\frac{1}{4}$ (4) $\frac{1}{5}$

PSC052

53. A collision is said to be perfectly inelastic when :-

- (1) Coefficient of restitution = 0
 (2) Coefficient of restitution = 1
 (3) Coefficient of restitution = ∞
 (4) Coefficient of restitution < 1

PSC053

54. A particle falls from a height 'h' upon a fixed horizontal plane and rebounds. If 'e' is the coefficient of restitution the total distance travelled before rebounding has stopped is :-

- (1) $h \left(\frac{1+e^2}{1-e^2} \right)$ (2) $h \left(\frac{1-e^2}{1+e^2} \right)$
 (3) $\frac{h}{2} \left(\frac{1-e^2}{1+e^2} \right)$ (4) $\frac{h}{2} \left(\frac{1+e^2}{1-e^2} \right)$

PSC054

55. If two masses m_1 and m_2 collide, the ratio of the changes in their respective velocities is proportional to :-

- (1) $\frac{m_1}{m_2}$ (2) $\sqrt{\frac{m_1}{m_2}}$ (3) $\frac{m_2}{m_1}$ (4) $\sqrt{\frac{m_2}{m_1}}$


PSC055

56. Two particles of mass M_A and M_B and their velocities are V_A and V_B respectively collide. After collision they interchange their velocities then ratio of $\frac{M_A}{M_B}$ is :-

- (1) $\frac{V_A}{V_B}$ (2) $\frac{V_B}{V_A}$
 (3) $\frac{(V_A + V_B)}{(V_B - V_A)}$ (4) 1

PSC056

57. In the diagrams given below the horizontal line represents the path of a ball coming from left and hitting another ball which is initially at rest. The other two lines represent the paths of the two balls after the collision. Which of the diagram shows a physically impossible situation?

- (1)  (2) 
 (3)  (4) 

PSC057

58. Two identical balls, one moves with 12 m/s and second is at rest, collide elastically. After collision velocity of second and first ball will be :-

- (1) 6m/s, 6m/s (2) 12m/s, 12m/s
 (3) 12m/s, 0m/s (4) 0m/s, 12m/s

PSC058

59. A sphere P of mass m and velocity $\vec{v}_i$ undergoes an oblique and perfectly elastic collision with an identical sphere Q initially at rest. The angle θ between the velocities of the spheres after the collision shall be :-

- (1) 0 (2) 45° (3) 90° (4) 180°

PSC059

60. A ball is dropped from a height of 10 m. If 40% of its energy is lost on collision with the earth then after collision the ball will rebound to a height of-

- (1) 10 m (2) 8 m (3) 4 m (4) 6 m

PSC060

61. A rubber ball is dropped from a height of 5m on a plane, where the acceleration due to gravity is not shown. On bouncing it rises to 1.8 m. The ball loses its velocity on bouncing by a factor of :-

- (1) $\frac{16}{25}$ (2) $\frac{2}{5}$ (3) $\frac{3}{5}$ (4) $\frac{9}{25}$

PSC061

62. Which of the following is true :-

- (1) Momentum is conserved in all collisions but kinetic energy is conserved only in inelastic collision
 (2) Neither momentum nor kinetic energy is conserved in inelastic collisions.
 (3) Momentum is conserved in all collisions but not kinetic energy
 (4) Both momentum and kinetic energy are conserved in all collisions.

PSC062

63. A bullet of mass m is fired into a large block of wood of mass M with velocity v. The final velocity of the system is :-

- (1) $\left(\frac{m}{M-m} \right) v$ (2) $\left(\frac{m+M}{M} \right) v$
 (3) $\left(\frac{M-m}{M} \right) v$ (4) $\left(\frac{m}{m+M} \right) v$

PSC063

Collisions and Centre of Mass

64. A big ball of mass M , moving with velocity u strikes a small ball of mass m , which is at rest. Finally small ball attains velocity u and big ball v . What is the value of v :-

(1) $\frac{M-m}{M}u$ (2) $\frac{m}{M+m}u$
 (3) $\frac{2m}{M+m}$ (4) $\frac{M}{M+m}v$

PSC064

65. A particle of mass m moving with speed v towards east strikes another particle of same mass moving with same speed v towards north. After striking, the two particles fuse together. With what speed this new particle of mass $2m$ will move in north-east direction?

(1) v (2) $\frac{v}{2}$ (3) $\frac{v}{\sqrt{2}}$ (4) $v\sqrt{2}$

PSC065

66. Two ice skaters A and B approach each other at right angles. Skater A has a mass 30 kg and velocity 1 m/s and skater B has a mass 20 kg and velocity 2 m/s . They meet and cling together. Their final velocity of the couple is

(1) 2 m/s
 (2) 1.5 m/s
 (3) 1 m/s
 (4) 2.5 m/s

PSC066

EXERCISE-I (Conceptual Questions)

ANSWER KEY

Question	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15
Answer	2	4	3	2	3	2	4	2	2	3	1	1	1	2	4
Question	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30
Answer	3	3	3	2	2	4	1	1	1	3	4	3	3	3	3
Question	31	32	33	34	35	36	37	38	39	40	41	42	43	44	45
Answer	2	1	3	4	2	1	4	3	4	1	2	2	2	3	2
Question	46	47	48	49	50	51	52	53	54	55	56	57	58	59	60
Answer	3	2	3	2	2	3	2	1	1	3	4	3	3	3	4
Question	61	62	63	64	65	66									
Answer	2	3	4	1	3	3									

Exercise - II (Previous Year Questions)

AIPMT/NEET

AIPMT 2015

1. Two particles of masses m_1 , m_2 move with initial velocities u_1 and u_2 . On collision, one of the particles get excited to higher level, after absorbing energy ε . If final velocities of particles be v_1 and v_2 then we must have :

$$(1) \frac{1}{2}m_1u_1^2 + \frac{1}{2}m_2u_2^2 = \frac{1}{2}m_1v_1^2 + \frac{1}{2}m_2v_2^2 - \varepsilon$$

$$(2) \frac{1}{2}m_1u_1^2 + \frac{1}{2}m_2u_2^2 - \varepsilon = \frac{1}{2}m_1v_1^2 + \frac{1}{2}m_2v_2^2$$

$$(3) \frac{1}{2}m_1^2u_1^2 + \frac{1}{2}m_2^2u_2^2 + \varepsilon = \frac{1}{2}m_1^2v_1^2 + \frac{1}{2}m_2^2v_2^2$$

$$(4) m_1^2u_1 + m_2^2u_2 - \varepsilon = m_1^2v_1 + m_2^2v_2$$

PSC074

2. Two spherical bodies of mass M and $5M$ and radii R and $2R$ are released in free space with initial separation between their centres equal to $12R$. If they attract each other due to gravitational force only, then the distance covered by the smaller body before collision is :-
(1) $4.5R$ (2) $7.5R$ (3) $1.5R$ (4) $2.5R$

PSC075

Re-AIPMT 2015

3. A ball is thrown vertically downwards from a height of 20 m with an initial velocity v_0 . It collides with the ground, loses 50% of its energy in collision and rebounds to the same height. The initial velocity v_0 is :

(Take $g = 10\text{ m/s}^2$)

$$(1) 10\text{ m/s} \quad (2) 14\text{ m/s}$$

$$(3) 20\text{ m/s} \quad (4) 28\text{ m/s}$$

PSC076

4. On a frictionless surface, a block of mass M moving at speed v collides elastically with another block of same mass M which is initially at rest. After collision the first block moves at an angle θ to its initial direction and has a speed $\frac{v}{3}$. The second block's speed after the collision is :-

$$(1) \frac{\sqrt{3}}{2}v \quad (2) \frac{2\sqrt{2}}{3}v \quad (3) \frac{3}{4}v \quad (4) \frac{3}{\sqrt{2}}v$$

PSC077

NEET-II 2016

5. A bullet of mass 10 g moving horizontally with a velocity of 400 m/s strikes a wooden block of mass 2 kg which is suspended by a light inextensible string of length 5 m . As a result, the centre of gravity of the block is found to rise a vertical distance of 10 cm . The speed of the bullet after it emerges out horizontally from the block will be :-

$$(1) 120\text{ m/s} \quad (2) 160\text{ m/s}$$

$$(3) 100\text{ m/s} \quad (4) 80\text{ m/s}$$

PSC078

6. Two identical balls A and B having velocities of 0.5 m/s and -0.3 m/s respectively collide elastically in one dimension. The velocities of B and A after the collision respectively will be :-

$$(1) -0.3\text{ m/s} \text{ and } 0.5\text{ m/s}$$

$$(2) 0.3\text{ m/s} \text{ and } 0.5\text{ m/s}$$

$$(3) -0.5\text{ m/s} \text{ and } 0.3\text{ m/s}$$

$$(4) 0.5\text{ m/s} \text{ and } -0.3\text{ m/s}$$

PSC079

NEET(UG) 2018

7. A moving block having mass m , collides with another stationary block having mass $4m$. The lighter block comes to rest after collision. When the initial velocity of the lighter block is v , then the value of coefficient of restitution (e) will be :-

$$(1) 0.5 \quad (2) 0.25 \quad (3) 0.8 \quad (4) 0.4$$

PSC080

NEET(UG) 2019

8. Body A of mass $4m$ moving with speed u collides with another body B of mass $2m$, at rest. The collision is head on and elastic in nature. After the collision the fraction of energy lost by the colliding body A is :

$$(1) \frac{1}{9} \quad (2) \frac{8}{9}$$

$$(3) \frac{4}{9} \quad (4) \frac{5}{9}$$

PSC081

NEET(UG) 2019 (Odisha)

9. An object flying in air with velocity $(20\hat{i} + 25\hat{j} - 12\hat{k})$ suddenly breaks in two pieces whose masses are in the ratio 1 : 5. The smaller mass flies off with a velocity $(100\hat{i} + 35\hat{j} + 8\hat{k})$. The velocity of the larger piece will be :-

- (1) $4\hat{i} + 23\hat{j} - 16\hat{k}$ (2) $-100\hat{i} - 35\hat{j} - 8\hat{k}$
 (3) $20\hat{i} + 15\hat{j} - 80\hat{k}$ (4) $-20\hat{i} - 15\hat{j} - 80\hat{k}$

PSC082

10. A particle of mass 5 m at rest suddenly breaks on its own into three fragments. Two fragments of mass m each move along mutually perpendicular direction with speed v each. The energy released during the process is :

- (1) $\frac{3}{5}mv^2$ (2) $\frac{5}{3}mv^2$
 (3) $\frac{3}{2}mv^2$ (4) $\frac{4}{3}mv^2$

PSC083

NEET(UG) 2020


11. Two particles of mass 5 kg and 10 kg respectively are attached to the two ends of a rigid rod of length 1 m with negligible mass. The centre of mass of the system from the 5 kg particle is nearly at a distance of :

- (1) 80 cm (2) 33 cm
 (3) 50 cm (4) 67 cm

PSC084

NEET(UG) 2020 (COVID-19)

12. Three identical spheres, each of mass M, are placed at the corners of a right angle triangle with mutually perpendicular sides equal to 2 m (see figure). Taking the point of intersection of the two mutually perpendicular sides as the origin, find the position vector of centre of mass.



- (1) $2(\hat{i} + \hat{j})$ (2) $(\hat{i} + \hat{j})$
 (3) $\frac{2}{3}(\hat{i} + \hat{j})$ (4) $\frac{4}{3}(\hat{i} + \hat{j})$

PSC085

NEET(UG) 2022

13. Two objects of mass 10 kg and 20 kg respectively are connected to the two ends of a rigid rod of length 10 m with negligible mass. The distance of the center of mass of the system from the 10 kg mass is :

- (1) $\frac{20}{3}$ m (2) 10 m
 (3) 5 m (4) $\frac{10}{3}$ m

PSC086

14. A shell of mass m is at rest initially. It explodes into three fragments having mass in the ratio 2 : 2 : 1. If the fragments having equal mass fly off along mutually perpendicular directions with speed v, the speed of the third (lighter) fragment is

- (1) $\sqrt{2}v$ (2) $2\sqrt{2}v$
 (3) $3\sqrt{2}v$ (4) v

PSC087

RE-NEET(UG) 2022

15. Given below are two statements : one is labelled as Assertion (A) and the other is labelled as Reason (R),

Assertion (A) :

When a fire cracker (rocket) explodes in mid air, its fragments fly in such a way that they continue moving in the same path, which the fire cracker would have followed, had it not exploded.

Reason (R) :

Explosion of cracker (rocket) occurs due to internal forces only and no external force acts for this explosion.

In the light of the above statements, choose the most appropriate answer from the options given below :

- (1) Both (A) and (R) are correct and (R) is the correct explanation of (A)
 (2) Both (A) and (R) are correct but (R) is not the correct explanation of (A)
 (3) (A) is correct but (R) is not correct
 (4) (A) is not correct but (R) is correct

PSC088

RE-NEET(UG) 2023		NEET(UG) 2024
16. A bullet of mass m hits a block of mass M elastically. The transfer of energy is the maximum, when : (1) $M = m$ (2) $M = 2m$ (3) $M \ll m$ (4) $M \gg m$	PSC097	18. Two bodies A and B of same mass undergo completely inelastic one-dimensional collision. The body A moves with velocity v_1 while body B is at rest before collision. The velocity of the system after collision is v_2 . The ratio $v_1 : v_2$ is : (1) 1 : 2 (2) 2 : 1 (3) 4 : 1 (4) 1 : 4
	PSC098	
17. Two particles A and B initially at rest, move towards each other under mutual force of attraction. At an instance when the speed of A is v and speed of B is $3v$, the speed of centre of mass is : (1) $2v$ (2) zero (3) v (4) $4v$	PSC099	

EXERCISE-II (Previous Year Questions)													ANSWER KEY		
Question	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15
Answer	2	2	3	2	1	4	2	2	1	4	4	3	1	2	4
Question	16	17	18												
Answer	1	2	2												

Exercise - III (Analytical Questions)

Master Your Understanding

1. Given below are two statements: One is labelled as **Assertion (A)** and the other is labelled as **Reason (R)**.

Assertion (A) : Two blocks of masses m_1 & m_2 are at rest. They are now moving towards each other under a mutual internal force. The velocity of centre of mass is zero.

Reason (R) : If no external force acts on the system, then velocity of centre of mass remains unchanged but can never be zero. In the light of the above statements, choose the most appropriate answer from the options given below:

- (1) Both **(A)** and **(R)** are true and **(R)** is the correct explanation of **(A)**.
 (2) Both **(A)** and **(R)** are true and **(R)** is NOT the correct explanation of **(A)**.
 (3) **(A)** is true but **(R)** is false.
 (4) **(A)** is false but **(R)** is true.

PSC089

2. Which of following statement is true.

- (1) COM of a rigid body is always inside the body.
 (2) Mass must be present physically at COM.
 (3) COM of two particle system is closer to particle with greater mass.
 (4) All above statements are true.

PSC091

3. Which of following statement is false.

- (1) KE in an elastic collision between two bodies remains same before and after collision.
 (2) There may or may not be physical contact between two bodies for collision to occur.
 (3) KE of two object system remains conserved during collision in elastic collision.
 (4) None of the above

PSC092

4. A projectile is projected at a speed u at an angle θ with horizontal. At the highest point projectile splits into two fragments of mass ratio 1 : 2. The smaller fragment coming to rest. Then consider following statements :-




- (A) In a given time both parts will cover equal vertical displacement.
 (B) COM of two fragments will follow the same path as the object would have followed if not exploded.
 (C) Linear momentum is conserved in vertical direction only.

Which of the following is correct?

- (1) A, B, C (2) A, B (3) B, C (4) A, C

PSC093

5. Match the following columns with respective positions of centre of mass -

Column - I		Column - II	
(A)	 uniform rod length = R	(P)	$x = \frac{3R}{8}$
(B)	 uniform thin semicircular ring of radius R	(Q)	$x = \frac{R}{2}$
(C)	 uniform semicircular disc of radius R	(R)	$x = \frac{4R}{3\pi}$
(D)	 uniform solid hemisphere of radius R	(S)	$x = \frac{2R}{\pi}$

- (1) A : P, B : R, C : Q, D : S
 (2) A : Q, B : P, C : R, D : S
 (3) A : Q, B : S, C : R, D : P
 (4) A : S, B : P, C : Q, D : R

PSC095

EXERCISE-III (Analytical Questions)

ANSWER KEY

Question	1	2	3	4	5
Answer	3	3	3	2	3

Important Notes